

# Nonflatness and totality

Basil A. Karádis

DRAFT OF 8 FEB 2018, 12:53 P.M.

## Abstract

We interpret finite types as domains over nonflat inductive base types in order to bring out the finitary core that seems to be inherent in the concept of totality. We prove a strong version of the Kreisel density theorem by providing a total compact element as a witness, a result that we cannot hope to have if we work with flat base types. To this end we develop tools that deal adequately with possibly inconsistent finite sets of information. The classical density theorem is reestablished via a “finite density theorem”, and corollaries are obtained, among them Berger’s separation property.

## 1 Introduction

In the area of denotational semantics of functional programming it is standard to view data types as countably based Scott domains in the tradition that started with Dana Scott’s and Yuri Ershov’s independent work in the late sixties and early seventies. More particularly, we may view these domains through their representations as Scott information systems, where programs are representatives of typed terms  $x : \rho$  with denotations being ideals in appropriate information systems, that is, consistent and deductively closed sets of tokens  $a \in x$ ; ideals are approximated by finite sets  $U \subseteq x$ , their so called formal neighborhoods.

A crucial choice in our setting is to work with *nonflat* rather than flat domains for the base types. These arise when we model base type partiality not as an extra pseudotoken, but as an extra nullary pseudoconstructor, which participates in the formation of further tokens and therefore leads to varying degrees of partiality. For example, while the *flat natural numbers*  $\{\perp, 0, S0, SS0, \dots\}$  feature just the bottom element for partiality, the *nonflat natural numbers* (also called *lazy natural numbers*)  $\{\perp, 0, S\perp, S0, SS\perp, SS0, \dots\}$  feature several partial elements, like  $\perp, S\perp, SS\perp$ , and so on; elements that do not involve  $\perp$ , like  $0, S0$ , or  $SS0$ , are called *total*. A basic advantage of this feature compared to flat base types is that we obtain injectivity and disjoint ranges for the constructors.

More generally, base-type nonflatness yields domains which are in a certain sense both richer, in that they contain more tokens, and tidier, in that they are finitely branching. Such domains seem to accommodate arguments that a flat setting cannot afford, and this paper intends to give one nontrivial example of this kind: an explicitly finitary approach to the *Kreisel density theorem*, a key result in the theory of higher-type computability.

Density was first stated and proved by Georg Kreisel in [25], and in different terms by Stephen Kleene in [24]. Building on work of Yuri Ershov [12, 13, 14], Ulrich Berger [2, 3] generalized and established density within domain theory, drawing as a corollary that it holds for the hierarchy of the partial continuous functionals over all finite types, and thus recovering the Kleene–Kreisel continuous functionals as equivalence

classes of the abstractly total elements in the hierarchy. Helmut Schwichtenberg and collaborators have already carried Berger's argument from the top (abstract domains with totality) down to the bottom (concrete Scott information systems induced by algebra constructors) numerous times in the past, starting with [42] and following up with [43, 19, 20, 44]. The present work is related to these top-down approaches, but attacks the problem in the opposite, bottom-up manner: we don't merely adapt previous domain-theoretic proofs to the setting of approximations, but rather work our way up, from approximations to ideals.

Let us recall the content of the Kreisel density theorem. We work with finite types, that is, with base types like  $\mathbb{N}$  and  $\mathbb{B}$  for naturals and booleans respectively, and then with arrow types above them. We capture the concept of termination by a totality predicate  $G$ :<sup>1</sup> at base types  $t$ , an ideal  $x$  is total if it contains a total token; at type  $\rho \rightarrow \sigma$ , an ideal  $f$  is total when it preserves totality, that is, when

$$\forall_{x:\rho} (G_\rho(x) \rightarrow G_\sigma(fx)) ,$$

where  $b \in fx$  for a token  $b \in \text{Tok}_\sigma$  if and only if  $\langle U, b \rangle \in f$ , for some formal neighborhood  $U \in \text{Con}_\rho$  with  $U \subseteq x$ . The *density property* for a type, the latter being understood as a space governed by the Scott topology, alleviates the omnipresent partiality by ensuring that every open set in the space nurtures total points, in other words, that total points are dense in the space. We formulate this here by saying that  $\rho$  is *dense* when

$$\forall_{U \in \text{Con}_\rho} \exists_{x:\rho} (G_\rho(x) \wedge U \subseteq x) . \quad (\text{D})$$

The Kreisel density theorem says that *every type is dense*.

Let us look a bit closer at the statement (D). We are given a neighborhood  $U$ , morally a compact element, comprising *finite* information, and we are supposed to come up with an ideal  $x$  as a witness, which may in principle be an *infinite* set of tokens. It is reasonable to suspect that the element of infinity in  $x$  must be inessential as far as an actual process of "totalization" of  $U$  is concerned—whatever this process might be—and that there is nothing inherently infinitary about it, since totalizing  $U$  should depend on its finite information, and not on the fact that the resulting ideal may be infinite. Can we then devise a totalization process which will feature an explicitly finitary core, that is,

*can we provide a witness for density which will be obviously finitary?*

We can easily see that in general we cannot do this if we interpret our base types by flat domains: At type  $\mathbb{N} \rightarrow \mathbb{N}$ , consider the very simple compact given intuitively by  $\{0 \mapsto \perp\}$ ; extending it, say, to  $\{0 \mapsto 0\}$  is not making it total, since it cannot respond (with a total value) to any total input different than 0; if we want to extend it to a total element, we must account for every possible total input except 0, which inevitably leads to an infinite set, say to the set  $\{0 \mapsto 0\} \cup \{S^m 0 \mapsto S0 \mid m > 0\}$ . This totalization process consists of two steps, namely totalizing the output on input zero from  $\perp$  to 0, and setting all other outputs to be  $S0$ ; nevertheless, the witness can only be presented as an infinite set. On the other hand, if we had interpreted the natural numbers by their nonflat domain, then we could have extended the compact  $\{0 \mapsto \perp\}$  to the compact

<sup>1</sup>The letter "g"—which probably derives from the English word *general* (for *generally defined*), and perhaps is influenced by the German word *gesamt* (complete, total, whole)—was used in this context already by Ershov [12, 14]. We adopt this notation here to designate totality in order to avoid confusion with other terms beginning with "t" in our text.

$\{0 \mapsto 0, S\perp \mapsto S0\}$ , which would have sufficed, since the information  $S\perp$  is enough to accommodate (later we will say "accept") all total numbers different than zero.

Our strategy can be summarized as follows. Step 1: define an appropriate notion of "total neighborhood". Step 2: establish a "finite density theorem", that is, that every neighborhood extends to such a total neighborhood. Step 3: show that a total neighborhood extends to a total ideal in a straightforward way.

We begin in section 2 with a necessary preamble on domains over nonflat base types represented by information systems. In section 3 we prepare for Step 1. As we can already see in the example that we gave above, one thing we have to do in order to "totalize" a neighborhood is to appropriately extend the set of its inputs; this finite set is seldom consistent, and in order to argue rigorously about it we must first develop a general understanding of such sets and their intricacies. This section delays the exposition of our argument a bit, but, beside gathering necessary definitions and facts, it will hopefully help familiarize the reader with these intricacies. Alternatively, the reader could skip to the next section and come back when the need arises. In section 4 we perform Steps 1 and 2: we define finite totality and prove finite density with Theorem 4.7; moreover, we characterize finite totality in a noninductive way in Theorem 4.9. Section 5 is about Step 3: with Theorem 5.10 we show how to obtain the classical Kreisel density theorem from the finite density theorem; in addition, we list some direct consequences, among them Berger's "separation property" in Proposition 5.14. We end in section 6 with comments on the literature and future work.

## 2 Nonflat domains via coherent information systems

We concentrate on a type system supporting arrow types over inductive base types.<sup>2</sup> We use  $\xi$  as a dummy type variable. Write  $\vec{\rho} \rightarrow \sigma$  to mean  $\rho_1 \rightarrow \dots \rightarrow \rho_r \rightarrow \sigma$  for some  $r \geq 0$  associated to the right; in case  $r = 0$  the vector is empty.

- For every vector  $\vec{\xi}$  of length  $r$ , the expression  $\vec{\xi} \rightarrow \xi$  is a *constructor type* (of arity  $r$ ).
- If  $\kappa_1, \dots, \kappa_k$  are constructor types for  $k > 0$  and one of them nullary, then  $\mu_\xi(\kappa_1, \dots, \kappa_k)$  is a *type*. We think of such types as inductively defined *base types* or *algebras*, generated by *constructors*  $C_l$  corresponding to  $\kappa_l$ , for  $l = 1, \dots, k$ .
- If  $\rho, \sigma$  are types then  $\rho \rightarrow \sigma$  is a *type*; these are the usual *higher types*.

Note that constructor types only serve to build base types, and are not themselves admitted as types. Examples of base types are

- the *unit type*  $\mathbb{U} := \mu_\xi(\xi)$  with a single nullary constructor,
- the *type of boolean values*  $\mathbb{B} := \mu_\xi(\xi, \xi)$ , with constructors for the truth  $\text{tt} : \mathbb{B}$  and the falsity  $\text{ff} : \mathbb{B}$ ,
- the *type of natural numbers*  $\mathbb{N} := \mu_\xi(\xi, \xi \rightarrow \xi)$ , with constructors for the zero  $0 : \mathbb{N}$  and the successor  $S : \mathbb{N} \rightarrow \mathbb{N}$ ,

<sup>2</sup>In this section we omit proofs and details, for which the reader may consult [44, Part 3] and [46, Part I]. In relation to the former, in particular, note that we will be working within the *nonparametric* and *finitary* fragment of the system.

- the *type of (extended) derivations*  $\mathbb{D} := \mu_{\xi}(\xi, \xi, \xi \rightarrow \xi, \xi \rightarrow \xi \rightarrow \xi)$ , with constructors for an axiom  $0 : \mathbb{D}$ , another axiom  $1 : \mathbb{D}$ , a one-premise rule  $S : \mathbb{D} \rightarrow \mathbb{D}$ , and a two-premise rule  $B : \mathbb{D} \rightarrow \mathbb{D} \rightarrow \mathbb{D}$  (this algebra is simple yet nontrivial enough to provide us with examples as we go along).

We will write  $t$  to denote an arbitrary *base* type and  $\rho, \sigma$  to denote arbitrary types in general.

A (*Scott*) *information system* [45, 47] is a triple  $(\text{Tok}, \text{Con}, \vdash)$ , where  $\text{Tok}$  is an inhabited countable set of *tokens*,  $\text{Con}$  is a collection of finite sets of tokens which we call *consistent sets* or (*formal*) *neighborhoods*, and  $\vdash$  is a subset of  $\text{Con} \times \text{Tok}$ , the *entailment*. These are subject to the axioms

$$\begin{aligned} \{a\} &\in \text{Con}, \\ U \subseteq V \wedge V \in \text{Con} &\rightarrow U \in \text{Con}, \\ U \in \text{Con} \wedge a \in U &\rightarrow U \vdash a, \\ U \vdash V \wedge V \vdash c &\rightarrow U \vdash c, \\ U \in \text{Con} \wedge U \vdash b &\rightarrow U \cup \{b\} \in \text{Con}, \end{aligned}$$

where  $U \vdash V$  stands for  $U \vdash b$  for all  $b \in V$ . From the latter follows vacuously that  $U \vdash \emptyset$  for all  $U$ , while  $\emptyset \in \text{Con}$  follows from the first two axioms. We may refer to the fifth axiom as *propagation (of consistency through entailment)*; note that this axiom may be equivalently expressed as

$$U \vdash V \rightarrow U \cup V \in \text{Con}. \quad (1)$$

For finite sets of tokens  $\Gamma$  which are not necessarily consistent we write  $\text{Fin}$ , so  $\text{Con} \subseteq \text{Fin}$ . An information system is called *coherent* when in addition to the above it satisfies

$$\forall_{a, a' \in U} \{a, a'\} \in \text{Con} \rightarrow U \in \text{Con} \quad (2)$$

for all  $U \in \text{Fin}$ . By the coherence and the second axiom above, it follows that the consistency of a token set is equivalent to the consistency of its pairs. Drawing on this property, we often write  $a \asymp b$  for  $\{a, b\} \in \text{Con}$ , and even  $U \asymp V$  for  $U \cup V \in \text{Con}$  (which is also often written  $U \uparrow V$ ). In the following we work exclusively with coherent systems, even if we don't mention it explicitly.

Given two coherent information systems  $\rho$  and  $\sigma$ , we form their *function space*  $\rho \rightarrow \sigma$ : define its tokens by  $\langle U, b \rangle \in \text{Tok}$  if  $U \in \text{Con}_{\rho}$  and  $b \in \text{Tok}_{\sigma}$ , its consistency by  $\langle U, b \rangle \asymp \langle U', b' \rangle$  if  $U \asymp_{\rho} U'$  implies  $b \asymp_{\sigma} b'$ , and its entailment by  $W \vdash \langle U, b \rangle$  if  $WU \vdash_{\sigma} b$ , where

$$b \in WU := \exists_{U' \in \text{Con}_{\rho}} (\langle U', b \rangle \in W \wedge U \vdash_{\rho} U').$$

The last operation is called *neighborhood application*. We will revisit it in some depth in section 3.3 where we will also show that it is monotone in both arguments, that is, that  $U \vdash U'$  implies  $WU \vdash WU'$  and that  $W \vdash W'$  implies  $WU \vdash WU'$  for all appropriate  $U, U', W, W'$  (Lemma 3.8). For the proof of the following see [44, Chapter 6].

**Fact 2.1.** *The function space of two coherent systems is itself a coherent information system.*

An *ideal* (or *element*) of an information system  $\rho$  is a possibly infinite token set  $x \subseteq \text{Tok}$ , such that  $U \in \text{Con}$  for every  $U \subseteq_f x$  (consistency), and  $U \vdash b$  for some  $U \subseteq_f x$  implies  $b \in x$  (deductive closure). If  $x$  is an ideal of  $\rho$ , we write  $x : \rho$  or  $x \in \text{Ide}_\rho$ . Note that in a generic setting built over flat base types, as for example the one described in [46], the empty set at every type  $\rho$  is an ideal, and plays the role of the bottom element  $\perp_\rho$ .

By a (*Scott–Ershov*) *domain* we mean here a countably based directed complete partial order with a least element, which is additionally algebraic and bounded complete. A domain is *coherent* [37], if every set of compacts has a least upper bound exactly when each of its *pairs* has a least upper bound. Write  $b \in \bar{U}$  if and only if  $U \vdash b$  (in the generic flat-based setting we have  $\bar{\emptyset} = \emptyset = \perp$ ). The following fact is directly based on the fundamental work of Scott [45]; for the proofs supporting our particular formulation see [46, Section 6.1] and [22, Theorem 8].

**Fact 2.2** (Representation theorem). *Let  $\rho = (\text{Tok}_\rho, \text{Con}_\rho, \vdash_\rho)$  be a coherent information system. Then  $(\text{Ide}_\rho, \subseteq, \perp_\rho)$  is a coherent domain with compacts given by  $\{\bar{U} \mid U \in \text{Con}_\rho\}$ . Conversely, every coherent domain can be represented by a coherent information system.*

An *approximable mapping* between two information systems  $\rho$  and  $\sigma$  is a relation  $r \subseteq \text{Con}_\rho \times \text{Con}_\sigma$  that generalizes entailment in the following sense:  $\langle \emptyset, \emptyset \rangle \in r$ ; if  $\langle U, V_1 \rangle, \langle U, V_2 \rangle \in r$  then  $V_1 \simeq_\sigma V_2$  and  $\langle U, V_1 \cup V_2 \rangle \in r$ ; and if  $U \vdash_\rho U'$ ,  $\langle U', V' \rangle \in r$ , and  $V' \vdash_\sigma V$ , then  $\langle U, V \rangle \in r$ . One can show [45] that there is a bijective correspondence between the approximable mappings from  $\rho$  to  $\sigma$  and the ideals of the function space  $\rho \rightarrow \sigma$ , and moreover establish the categorical equivalence between domains with Scott continuous functions and information systems with approximable mappings. The equivalence is preserved if we restrict ourselves to the coherent case on both sides [22].

The *Scott topology* on  $\text{Ide}_\rho$  is given by the collection  $\{\nabla U \mid U \in \text{Con}_\rho\}$ , where  $\nabla U$  is the set  $\{x : \rho \mid U \subseteq x\}$  of all ideals above  $U$ . A set  $\mathcal{U} \subseteq \text{Ide}_\rho$  of ideals is *Scott open* when it is closed under supersets (Alexandrov condition) and for every  $x \in \mathcal{U}$  there is a  $U \subseteq x$  such that  $\bar{U} \in \mathcal{U}$  (Scott condition). One can furthermore show that an ideal-mapping  $f$  sending ideals from  $\text{Ide}_\rho$  to ideals in  $\text{Ide}_\sigma$  is *Scott continuous* when it is *monotone* and satisfies the *principle of finite support* (also called *approximation principle*) for all  $x : \rho$ , that is,

$$\bigvee_{b \in \text{Tok}_\sigma} (b \in f(x) \rightarrow \exists_{U \in \text{Con}_\rho} (U \subseteq x \wedge b \in f(\bar{U}))). \quad (\text{FS})$$

Finally, it can be shown (see for example [44, §6.1.3]) that the ideals  $\text{Ide}_{\rho \rightarrow \sigma}$  and the Scott continuous ideal-mappings  $\text{Ide}_\rho \rightarrow \text{Ide}_\sigma$  are in a bijective correspondence, a fact that justifies the nondiscriminating notation  $f : \rho \rightarrow \sigma$ .

Now we proceed to assign an information system to each type. Every higher type is naturally assigned a function space, so it suffices to discuss the information systems for base types, that is, for algebras. Let  $t$  be an algebra, with at least one nullary constructor if it is to be nontrivial. We add to it an extra nullary pseudoconstructor  $*_t$  (or just  $*$ ) to denote partiality (which we nevertheless agree to not let it appear in the algebra signature). This is a departure from [44], where partiality is treated as a special *untyped* symbol which is used by all possible algebras; for us, every algebra has its own partiality symbol. We define  $\text{Tok}_t$ ,  $\text{Con}_t$ , and  $\vdash_t$  inductively.

- If  $C$  is an  $r$ -ary constructor and  $a_i \in \text{Tok}_t$  for  $i = 1, \dots, r$  then  $Ca_1 \cdots a_r \in \text{Tok}_t$ .<sup>3</sup>

<sup>3</sup>Throughout the text we adopt the polish notation for tokens for typographical convenience.

For its *head constructor* write  $\text{hd}(Ca_1 \cdots a_r) = C$ ; for its  *$i$ -th component token* write  $a(i)$ , that is,  $(Ca_1 \cdots a_r)(i) = a_i$  for  $i = 1, \dots, r$ .

- We have  $a \succsim_l *$  and  $* \succsim_l a$  for all  $a \in \text{Tok}_l$ . Furthermore, if  $C$  is an  $r$ -ary constructor and  $a_i \succsim_l b_i$  for  $i = 1, \dots, r$  then  $Ca_1 \cdots a_r \succsim_l Cb_1 \cdots b_r$ . Finally, we have  $U \in \text{Con}_l$  if  $a \succsim_l a'$  for all  $a, a' \in U$ .
- We have  $U \vdash_l *$  for all  $U \in \text{Con}_l$ . Furthermore, if  $C$  is an  $r$ -ary constructor, every  $U_i \in \text{Con}_l$  is inhabited and  $U_i \vdash_l b_i$  for  $i = 1, \dots, r$ , then  $U \vdash_l Cb_1 \cdots b_r$  for all  $U \in \text{Con}_l$  that are *sufficient for  $C$  on  $U_1, \dots, U_r$* , in the sense that for each  $i = 1, \dots, r$  and each  $a_i \in U_i$  there exists an  $a \in U$  such that  $\text{hd}(a) = C$  and  $a(i) = a_i$ . Finally, if  $U \vdash_l b$ , then also  $U \cup \{*\} \vdash_l b$ .

Here we need to be cautious. The definition of  $\text{Con}_l$  incorporates the coherence property (2), so it follows that  $\emptyset \vdash_l \{*\}$ . This seems to undermine Fact 2.2, since the empty set cannot be an ideal anymore. It is nevertheless straightforward to show that the representation theorem still stands if we reinterpret the bottom ideals as follows:

$$\begin{aligned} \perp_l &:= \{*\}, \\ \perp_{\rho \rightarrow \sigma} &:= \{\langle U, b \rangle \mid U \in \text{Con}_\rho \wedge b \in \perp_\sigma\}. \end{aligned}$$

It is maybe worth pointing out that while our situation diverges from the generic flat-based setting and even from settings like the one of [44], in the same time it resonates the original Scott axioms [45]: there exists at least one trivial token in every information system, in particular one at every base type and several at every higher type.

Concerning sufficiency, we note the following: (a) in case  $C$  is a proper constructor,  $U$  is sufficient for  $C$  on  $U_1, \dots, U_r$  if and only if  $U \cup \{*\}$  is, if and only if  $U \setminus \{*\}$  is, and (b) we trivially have  $U \vdash_l CU_1 \cdots U_r$ , whenever  $U$  is sufficient for  $C$  on  $U_1, \dots, U_r$ ; the *constructor application* here is defined by

$$CU_1 \cdots U_r := \{Ca_1 \cdots a_r \mid a_1 \in U_1, \dots, a_r \in U_r\},$$

which is consistent if and only if every  $U_i$  is consistent. Moreover, every neighborhood  $U$  that is nontrivial (meaning,  $\{*\} \not\vdash_l U$ ) is indeed equivalent to one of the form  $CU_1 \cdots U_r$ : if

$$U \setminus \{*\} = \{Ca_{11} \cdots a_{r1}, \dots, Ca_{1m} \cdots a_{rm}\},$$

we gather all  $i$ -th component tokens into a neighborhood, the  *$i$ -th component neighborhood*  $U(i) := \{a_{i1}, \dots, a_{im}\}$  of  $U$ , and let  $U_i := U(i)$  for every  $i = 1, \dots, r$ ; then we indeed have  $U \sim_l CU_1 \cdots U_r$  (where  $U \sim V$  abbreviates  $U \vdash V \wedge V \vdash U$ ).

The proof of the following is straightforward but tedious.

**Fact 2.3.** *Let  $\mathfrak{t}$  be an algebra given by constructors. The triple  $(\text{Tok}_l, \text{Con}_l, \vdash_l)$  is a coherent information system.*

### 3 Finite sets

Recall that the first step in our strategy is to decide on a reasonable definition of “finite totality”, one that will already embody the totalization mechanism for density on the one hand, and that will be susceptible to a canonical extension to a total ideal on the other. The nature of our study leads us naturally to define the concept by induction over types, which is what we do in section 4.1. Nevertheless, we will see in Theorem 4.9 that we can

arrive at the same concept in an explicit way in terms of “transitive elements”: elements that witness local transitivity within a not necessarily transitive relation—in our case, the consistency relation. To work with such sets of tokens we need an operation akin to application, defined using consistency rather than entailment. This is how we come to spend some space discussing not necessarily consistent finite sets in some generality, while we postpone the actual definition of total neighborhoods until section 4.

### 3.1 Entailment and consistency for finite sets

There is the trivial syntactical reason to look at finite sets in general and not just at the consistent ones: the latter presuppose the former by definition—in particular, the thematization of finite sets is unavoidable in implementation endeavors like [20]. But finite sets may play a natural and important role within purely semantical arguments as well—to mention a naive example, think of the subtokens  $a_1, \dots, a_r$  of a base-type token  $a = Ca_1 \cdots a_r$ . In this subsection we will hardly cover anything more than what we will need later, with the exception of Lemma 3.2, which we included for the sake of some points in §3.3.

As we already mentioned, we write  $\text{Fin}_\rho$  instead of  $\mathcal{P}_f(\text{Tok}_\rho)$ , so  $\Gamma \in \text{Fin}_\rho$  means that  $\Gamma$  is a *finite* set of tokens, not necessarily consistent. If  $\Theta = \{\langle U_j, b_j \rangle \mid j = 1, \dots, l\} \in \text{Fin}_{\rho \rightarrow \sigma}$ , write  $L(\Theta)$  for  $\bigcup_j U_j \in \text{Fin}_\rho$  (notice that this is a *flattening*), and  $R(\Theta)$  for  $\bigcup_j \{b_j\} \in \text{Fin}_\sigma$ . Furthermore, if  $U \in \text{Con}_\rho$  and  $\Delta \in \text{Fin}_\sigma$ , write  $\langle U, \Delta \rangle$  for  $\{\langle U, b \rangle \mid b \in \Delta\} \in \text{Fin}_{\rho \rightarrow \sigma}$  (note that  $\langle U, \emptyset_\sigma \rangle = \emptyset_{\rho \rightarrow \sigma}$ ).

**Lemma 3.1.** *Let  $\Theta, \Theta' \in \text{Fin}_{\rho \rightarrow \sigma}$ . We have  $L(\Theta \cup \Theta') = L(\Theta) \cup L(\Theta')$  and  $R(\Theta \cup \Theta') = R(\Theta) \cup R(\Theta')$ . Furthermore, if  $\Theta \subseteq \Theta'$  then we have  $L(\Theta) \subseteq L(\Theta')$  as well as  $R(\Theta) \subseteq R(\Theta')$ .*

A *neighborhood* in  $\Gamma \in \text{Fin}_\rho$  is a subset  $U \subseteq \Gamma$ , which happens to be consistent; write  $U \in \text{Con}_\Gamma$ . The empty set and the singletons of  $\Gamma$  are always in  $\text{Con}_\Gamma$ . Say that  $\Gamma$  *entails*  $\Gamma'$  (as a finite set), and write  $\Gamma \vdash_\rho^F \Gamma'$ , when

$$\forall U' \in \text{Con}_{\Gamma'} \exists U \in \text{Con}_\Gamma U \vdash_\rho U'.$$

This is a generalization of the notion  $U \vdash_\rho U'$  for neighborhoods. A simpler, but less helpful generalization is “ $\Gamma \vdash_\rho \Gamma'$  if and only if for every  $a' \in \Gamma'$  there is some  $U \in \text{Con}_\Gamma$  such that  $U \vdash_\rho a'$ ”; if  $\Gamma \vdash_\rho^F \Gamma'$  then also  $\Gamma \vdash_\rho \Gamma'$ . Contrary to the case of consistent sets, although  $\Gamma \vdash_\rho^F \Gamma'$  implies  $\Gamma \vdash_\rho^F a$  for all  $a \in \Gamma'$ , the converse is not true in general. For example  $\{B00, B11\} \vdash_D^F B0*$  and  $\{B00, B11\} \vdash_D^F B*1$ , but  $\{B00, B11\} \not\vdash_D^F \{B0*, B*1\}$ .

Similarly, say that  $\Gamma$  and  $\Gamma'$  are *consistent* (as finite sets), and write  $\Gamma \asymp_\rho^F \Gamma'$ , when

$$\forall U \in \text{Con}_\Gamma \forall U' \in \text{Con}_{\Gamma'} U \asymp_\rho U'.$$

Again, this is a generalization of consistency between neighborhoods which proves more useful for not necessarily consistent finite sets than the simpler notion “ $\Gamma \asymp_\rho \Gamma'$  if and only if  $\{a, a'\} \in \text{Con}_\rho$  for all  $a \in \Gamma$  and  $a' \in \Gamma'$ ” (which we may nevertheless occasionally use); if  $\Gamma \asymp_\rho^F \Gamma'$  then  $\Gamma \asymp_\rho \Gamma'$ . Note that in the case of  $\asymp_\rho^F$  we generally don’t have reflexivity; in fact, we trivially have  $\Gamma \asymp_\rho^F \Gamma$  if and only if  $\Gamma \in \text{Con}_\rho$ . An example of consistency between inconsistent finite sets is  $\{B0*, B1*\} \asymp_D^F \{B*0, B*1\}$ .

Reflexivity of consistency is the only property that the triple  $(\text{Fin}_\rho, \asymp_\rho^F, \vdash_\rho^F)$  lacks in order to constitute a Scott information system.

**Lemma 3.2.** *The entailment between finite sets is reflexive and transitive and the consistency between finite sets is symmetric and propagates through entailment, that is,*

1.  $\forall \Gamma \in \text{Fin} \Gamma \vdash^F \Gamma$ ,
2.  $\forall \Gamma, \Delta, \Theta \in \text{Fin} (\Gamma \vdash^F \Delta \wedge \Delta \vdash^F \Theta \rightarrow \Gamma \vdash^F \Theta)$ ,
3.  $\forall \Gamma, \Delta \in \text{Fin} (\Gamma \asymp^F \Delta \rightarrow \Delta \asymp^F \Gamma)$ ,
4.  $\forall \Gamma, \Delta, \Theta \in \text{Fin} (\Gamma \asymp^F \Delta \wedge \Delta \vdash^F \Theta \rightarrow \Gamma \asymp^F \Theta)$ .

*Proof.* We just show the propagation property. Let  $\Gamma, \Delta, \Theta \in \text{Fin}$  be such that  $\Gamma \asymp^F \Delta$  and  $\Delta \vdash^F \Theta$ . Consider  $U \in \text{Con}_\Gamma$  and  $W \in \text{Con}_\Theta$ ; by the assumptions, there exists a  $V \in \text{Con}_\Delta$  with  $V \vdash W$  and  $U \asymp V$ ; by propagation on  $\text{Con}$  (1) we get  $U \asymp W$ .  $\square$

### 3.2 Maximal and transitive neighborhoods

Think of some finite set  $\Gamma$  of type  $\rho$  and suppose that we wish to assign  $\sigma$ -values  $b_i$  to neighborhoods  $U_i$  of  $\text{Con}_\Gamma$  (for some  $i \in I$ ) in a way that the finite set  $\{\langle U_i, b_i \rangle \mid i \in I\}$  at type  $\rho \rightarrow \sigma$  will be consistent. Some reflection shows that it suffices to pair the “maximal” neighborhoods of  $\Gamma$  with the given values of  $\sigma$ , but we can actually do better than that: we can relax the requirement of maximality by requiring instead that we assign the given arbitrary values already to those neighborhoods which are, so to speak, maximal enough or “almost maximal”, in the sense that they are below exactly one maximal in  $\Gamma$ ; these are exactly the “transitive neighborhoods” in  $\Gamma$ .

Call  $U \in \text{Con}_\Gamma$  a *maximal neighborhood in  $\Gamma$* , and write  $U \in \text{Con}_\Gamma^{\text{max}}$ , when it is maximal with respect to the entailment relation, that is, when

$$\forall_{U' \in \text{Con}_\Gamma} (U' \vdash_\rho U \rightarrow U \vdash_\rho U').$$

Call  $U \in \text{Con}_\Gamma$  (*consistency*) *transitive in  $\Gamma$* , and write  $U \in \text{Con}_\Gamma^{\text{ctr}}$ , when it satisfies the property

$$\forall_{U_1, U_2 \in \text{Con}_\Gamma} (U_1 \asymp_\rho U \asymp_\rho U_2 \rightarrow U_1 \asymp_\rho U_2).$$

We can reformulate this by introducing the notation  $\tilde{U}$  for the *consistency closure* of  $U$ , that is, for the set  $\{a \in \text{Tok}_\rho \mid U \asymp_\rho a\}$  (it is clear that, while it encompasses the deductive closure, the consistency closure of a neighborhood is *not* in general an ideal, because consistency may fail); then  $U$  is transitive in  $\Gamma$  when  $\tilde{U} \cap \Gamma \in \text{Con}_\rho$ .

More generally, call  $U \in \text{Con}_\rho$  *transitive for  $\Gamma$*  or just  *$\Gamma$ -transitive (in  $\rho$ )*, and write  $U \in \text{Con}_{\rho|\Gamma}^{\text{ctr}}$ , if, again,  $U_1 \asymp_\rho U \asymp_\rho U_2$  implies  $U_1 \asymp_\rho U_2$  for all  $U_1, U_2 \in \text{Con}_\Gamma$ ; obviously,  $\text{Con}_\Gamma^{\text{ctr}} \subseteq \text{Con}_{\rho|\Gamma}^{\text{ctr}}$ .

It is clear that every maximal in a finite set is also transitive in it. It is also immediate that consistency between neighborhoods, restricted to  $\text{Con}_\Gamma^{\text{ctr}}$  (but not to  $\text{Con}_{\rho|\Gamma}^{\text{ctr}}$ !), becomes an equivalence relation. Still trivially, but importantly, we have the following.

**Lemma 3.3** (Upward closedness of transitivity). *Let  $\rho$  be a type and  $\Gamma \in \text{Fin}_\rho$ . For any  $U, U' \in \text{Con}_\rho$ , if  $U \in \text{Con}_{\rho|\Gamma}^{\text{ctr}}$  and  $U' \vdash_\rho U$ , then  $U' \in \text{Con}_{\rho|\Gamma}^{\text{ctr}}$ .*

*Proof.* Let  $U_1, U_2 \in \text{Con}_\Gamma$  be such that  $U_1 \asymp_\rho U' \asymp_\rho U_2$ . By propagation (1) we have  $U_1 \asymp_\rho U \asymp_\rho U_2$ , so  $U_1 \asymp_\rho U_2$ .  $\square$

It is often handy to check for extremality (that is, maximality or transitivity) on the level of tokens.

**Lemma 3.4** (Extremality through tokens). *Let  $\rho$  be an arbitrary type,  $\Gamma \in \text{Fin}_\rho$ , and  $U \in \text{Con}_\Gamma$ .*

1. *We have  $U \in \text{Con}_\Gamma^{\text{max}}$  if and only if  $U \asymp_\rho a$  implies  $U \vdash_\rho a$  for  $a \in \Gamma$ .*
2. *We have  $U \in \text{Con}_{\rho|\Gamma}^{\text{ctr}}$  if and only if  $a_1 \asymp_\rho U \asymp_\rho a_2$  implies  $a_1 \asymp_\rho a_2$  for  $a_1, a_2 \in \Gamma$ .*

*Proof.* For 1. From left to right, assume that  $U$  is maximal, and let  $a \in \Gamma$  be such that  $U \asymp_\rho a$ . Then  $U \cup \{a\} \vdash_\rho U$ , and by the maximality of  $U$  we get  $U \sim_\rho U \cup \{a\}$ , which gives us  $U \vdash_\rho a$ . For the other way around, let  $U' \in \text{Con}_\Gamma$  be such that  $U' \vdash_\rho U$ ; then  $U \asymp_\rho U'$ , and by the assumption we get that  $U \vdash_\rho U'$ , so  $U$  is indeed maximal.

For 2. From left to right, assume that  $U$  is transitive for  $\Gamma$ , and let  $a_1, a_2 \in \Gamma$  be such that  $U \asymp_\rho a_i$  for both  $i$ . Then  $U \asymp_\rho \{a_i\}$ , and by the transitivity of  $U$  we get  $a_1 \asymp_\rho a_2$ . For the other way around, let  $U_1, U_2 \in \text{Con}_\Gamma$  be such that  $U \asymp_\rho U_i$ , for both  $i$ , and  $a_i \in U_i$ ; then  $a_1 \asymp_\rho a_2$  by the assumption, so  $U_1 \asymp_\rho U_2$ , and  $U$  is indeed  $\Gamma$ -transitive.  $\square$

The lemma makes the significance of extremality in a finite set quite apparent. In particular, it is good to know that two maximals in a finite set are either equivalent or inconsistent (a fact that we can put even more bluntly like this: if  $U$  is maximal and deductively closed in  $\Gamma$ , then for each  $a \in \Gamma$  we have either  $a \in U$  or  $a \not\asymp_\rho U$ ).

**Lemma 3.5.** *Let  $\rho$  be any type. For all  $\Gamma \in \text{Fin}_\rho$  and  $U \in \text{Con}_\Gamma^{\text{max}}$ , if  $U' \in \text{Con}_\rho$  is such that  $U' \vdash_\rho U$  then  $U' \in \text{Con}_{\Gamma \cup U'}^{\text{max}}$ .*

*Proof.* Let  $a \in \Gamma \cup U'$  be such that  $a \asymp_\rho U'$ ; since  $U' \vdash_\rho U$ , we have  $a \asymp_\rho U$ . In case  $a \notin \Gamma$  we have  $a \in U'$ ; in case  $a \in \Gamma$ , we have  $U \vdash_\rho a$  by Lemma 3.4.1; in both cases it follows that  $U' \vdash_\rho a$ , so  $U'$  is maximal in  $\Gamma \cup U'$  by Lemma 3.4.1.  $\square$

**Lemma 3.6** (Maximal extensions). *Let  $\rho$  be a type and  $\Gamma \in \text{Fin}_\rho$ .*

1. *For any  $U \in \text{Con}_\Gamma$ , we have  $U \in \text{Con}_\Gamma^{\text{ctr}}$  if and only if there is exactly one  $\hat{U} \in \text{Con}_\Gamma^{\text{max}}$ , up to equivalentailment, such that  $\hat{U} \vdash_\rho U$ .*
2. *For any  $U \in \text{Con}_\rho$ , we have  $U \in \text{Con}_{\rho|\Gamma}^{\text{ctr}}$  if and only if, whenever there exist  $U_0 \in \text{Con}_\Gamma$  with  $U \asymp_\rho U_0$ , there exists a  $\hat{U} \in \text{Con}_\Gamma^{\text{max}}$  such that  $U \asymp_\rho U_0$  implies  $\hat{U} \vdash_\rho U_0$  for all  $U_0 \in \text{Con}_\Gamma$ .*

*Proof.* For 1, from left to right, assume that  $U \in \text{Con}_\Gamma^{\text{ctr}}$  and let  $U_1, U_2 \in \text{Con}_\Gamma^{\text{max}}$  be such that  $U_i \vdash_\rho U$  for both  $i = 1, 2$ . By the propagation of consistency (1), we have  $U_1 \asymp_\rho U \asymp_\rho U_2$ ; by the assumption we have  $U_1 \asymp_\rho U_2$ ; by the maximality of  $U_1$  and  $U_2$ , it follows from Lemma 3.4.1 that  $U_1 \sim_\rho U_2$ .

For the other direction, assume that  $U$  is such that any two maximal neighborhoods in  $\Gamma$  that entail it are equivalent, and let  $U_1, U_2 \in \text{Con}_\Gamma$  be such that  $U_1 \asymp_\rho U \asymp_\rho U_2$ . Then for any two  $U_1^m, U_2^m \in \text{Con}_\Gamma^{\text{max}}$ , with  $U_i^m \vdash_\rho U \cup U_i$ , by the assumption, we must have  $U_1^m \sim_\rho U_2^m$ ; it follows that  $U_1 \asymp_\rho U_2$ , by the propagation of consistency.

For 2, let  $U \in \text{Con}_\rho$ . Assume that  $U \in \text{Con}_{\rho|\Gamma}^{\text{ctr}}$  and  $U \asymp_\rho U_i$  for some  $U_i \in \text{Con}_\Gamma$ , where  $i > 0$ . Gather all these  $U_i$  in the neighborhood  $U_0 := \bigcup_i U_i$ ; we have of course  $U_0 \in \text{Con}_\Gamma$ . Then there is at least one maximal  $\hat{U} \in \text{Con}_\Gamma^{\text{max}}$  such that  $\hat{U} \vdash_\rho U_0 \vdash_\rho U_i$  for all  $i$ .

Conversely, assume that  $U$  satisfies

$$\exists_{U_0 \in \text{Con}_\Gamma} U \asymp_\rho U_0 \rightarrow \exists_{\hat{U} \in \text{Con}_\Gamma^{\text{max}}} \forall_{U_0 \in \text{Con}_\Gamma} (U \asymp_\rho U_0 \rightarrow \hat{U} \vdash_\rho U_0),$$

and let  $U_1, U_2 \in \text{Con}_\Gamma$  be such that  $U_1 \asymp_\rho U \asymp_\rho U_2$ . From the assumption we get a maximal  $\hat{U} \in \text{Con}_\Gamma^{\text{max}}$  with  $\hat{U} \vdash_\rho U_i$  for each  $i$ , so  $U_1 \asymp_\rho U_2$ .  $\square$

We will say *the maximal extension of  $U$  in  $\Gamma$* , if  $U \in \text{Con}_\Gamma^{\text{ctr}}$ , for the unique (up to equivalence) maximal neighborhood entailing  $U$ ; this we denote by  $\hat{U}$ , as in the statement of the above Lemma. But note that in the case of transitive neighborhoods *outside*  $\Gamma$  uniqueness is not guaranteed: an example with two maximals at type  $\mathbb{D}$  is provided by the finite set  $\Gamma = \{S^*, S0, S1\}$  and the neighborhood  $U = \{SS^*\}$ .

### 3.3 Upper and middle application

The notion of application  $fx$  of some higher-type term  $f$  to some input term  $x$ , both appropriately typed, is interpreted as “the information that we hold on  $x$  suffices to draw the information  $fx$  on the output, given the information that we have on  $f$ ”. In section 2, in the definition of neighborhood application, we saw that when we bring this notion down to the finite level it is *entailment* that we read into “suffices”, but for our purposes it will come in handy to consider a different version of application between neighborhoods, where we replace entailment by consistency. As will become clear in the results below, starting already with Lemma 3.10, this kind of application is justified by the existence and use of transitive neighborhoods.

Let  $\Theta \in \text{Fin}_{\rho \rightarrow \sigma}$  and  $U \in \text{Con}_\rho$ . The (*upper*) *application*  $\Theta \cdot U$  gathers all values  $b \in \text{Tok}_\sigma$  whose arguments  $U_b$  fall under  $U$ :

$$b \in \Theta \cdot U := \exists_{U_b \in \text{Con}_\rho} (\langle U_b, b \rangle \in \Theta \wedge U \vdash_\rho U_b).$$

Note that this trivially generalizes the neighborhood application of section 2; from now on we will always write  $W \cdot U$  instead of  $WU$ . The *middle application*  $\Theta \cdot U$  is defined by

$$b \in \Theta \cdot U := \exists_{U_b \in \text{Con}_\rho} (\langle U_b, b \rangle \in \Theta \wedge U \asymp_\rho U_b).$$

It follows immediately from the definition that the middle application yields at least as much information as the upper one does, namely  $\Theta \cdot U \subseteq \Theta \cdot U$ .

In the case of a consistent left argument, we can make the following easy observations.

**Lemma 3.7.** *Let  $\rho, \sigma$  be arbitrary types,  $W \in \text{Con}_{\rho \rightarrow \sigma}$ ,  $U \in \text{Con}_\rho$  and  $b \in \text{Tok}_\sigma$ .*

1. *We have  $W \vdash_{\rho \rightarrow \sigma} \langle U, b \rangle$  if and only if  $W \cdot U \vdash_\sigma b$ .*
2. *We have  $W \asymp_{\rho \rightarrow \sigma} \langle U, b \rangle$  if and only if  $W \cdot U \asymp_\sigma b$ .*

Note that in 3.7.2 the finite set  $W \cdot U$  may not be consistent, but we still did not write  $W \cdot U \asymp_\sigma^F b$ ; here we are just saying that every pair  $\{b^W, b\}$  will be consistent, for  $b^W \in W \cdot U$ .

**Lemma 3.8 (Application).** *Let  $\rho, \sigma$  be arbitrary types.*

1. *Application is consistently defined, that is, if  $W \in \text{Con}_{\rho \rightarrow \sigma}$  and  $U \in \text{Con}_\rho$ , then  $W \cdot U \in \text{Con}_\sigma$ .*
2. *Application is monotone in the right argument, in particular, if  $\Theta \in \text{Fin}_{\rho \rightarrow \sigma}$  and  $U, U' \in \text{Con}_\rho$ , with  $U \vdash_\rho U'$ , then  $\Theta \cdot U' \subseteq \Theta \cdot U$ .*
3. *Application is monotone in the left argument, that is, if  $\Theta, \Theta' \in \text{Fin}_{\rho \rightarrow \sigma}$  with  $\Theta \vdash_{\rho \rightarrow \sigma}^F \Theta'$  and  $U \in \text{Con}_\rho$ , then  $\Theta \cdot U \vdash_\sigma^F \Theta' \cdot U$ .*

*Proof.* For 1, let  $W \in \text{Con}_{\rho \rightarrow \sigma}$  and  $U \in \text{Con}_\rho$ , and consider  $b_1, b_2 \in W \cdot U$ . By the definition, there must exist  $\langle U_1, b_1 \rangle, \langle U_2, b_2 \rangle \in W$ , such that  $U \vdash_\rho U_1 \cup U_2$ ; it follows that  $U_1 \asymp_\rho U_2$ , so the consistency of  $W$  ensures that  $b_1 \asymp_\sigma b_2$ .

For 2, let  $\Theta \in \text{Fin}_{\rho \rightarrow \sigma}$  and  $U, U' \in \text{Con}_\rho$ , and assume that  $U \vdash_\rho U'$ . Consider a  $b \in V$ ; by the definition there exists a  $U_b \in L(\Theta)$  with  $\langle U_b, b \rangle \in \Theta$  and  $U' \vdash_\rho U_b$ ; the assumption immediately gives  $U \vdash_\rho U_b$ , so  $b \in \Theta \cdot U$  as well.

For 3, let  $\Theta, \Theta' \in \text{Con}_{\rho \rightarrow \sigma}$  and  $U \in \text{Con}_\rho$ , and assume that  $\Theta \vdash_{\rho \rightarrow \sigma}^F \Theta'$ . Consider a  $V' \in \text{Con}_{\Theta' \cdot U}$ ; for each  $b' \in V'$  there is a  $U_{b'} \in \text{Con}_\rho$  such that  $\langle U_{b'}, b' \rangle \in \Theta'$  and  $U \vdash_\rho U_{b'}$ ; the set  $W' := \{\langle U_{b'}, b' \rangle \in \Theta' \mid b' \in V' \wedge U \vdash_\rho U_{b'}\}$  is consistent in  $\Theta'$ . By the assumption there exists some  $W \in \text{Con}_\Theta$  such that  $W \vdash_{\rho \rightarrow \sigma} W'$ . Since for each  $\langle U_{b'}, b' \rangle \in W'$  we have  $W \cdot U_{b'} \vdash_\sigma b'$ , by 2 we get  $W \cdot U \vdash_\sigma W \cdot U_{b'}$ , hence  $W \cdot U \vdash_\sigma b'$ , that is,  $W \cdot U \vdash_\sigma V'$  and since  $W \cdot U \in \text{Con}_{\Theta \cdot U}$ , we are done.  $\square$

The following gives us conservative extensions of a neighborhood by way of extending its set of arguments.

**Lemma 3.9.** *Let  $W \in \text{Con}_{\rho \rightarrow \sigma}$  and  $\Gamma \in \text{Fin}_\rho$  such that  $L(W) \subseteq \Gamma$ . Then*

$$W \sim_{\rho \rightarrow \sigma} \bigcup_{U \in \text{Con}_\Gamma} \langle U, W \cdot U \rangle.$$

*Proof.* From left to right, let  $U \in \text{Con}_\Gamma$  and  $b \in W \cdot U$ . There exists a  $U_b \in \text{Con}_{L(\Theta)}$  with  $\langle U_b, b \rangle \in W$  and  $U \vdash_\rho U_b$ . Then  $\langle U_b, b \rangle \vdash_{\rho \rightarrow \sigma} \langle U, b \rangle$ . The other way around is obvious, since  $W \subseteq \bigcup_{U \in \text{Con}_\Gamma} \langle U, W \cdot U \rangle$ .<sup>4</sup>  $\square$

Turning our attention to middle application, the first thing we want to know is how it fares compared to Lemma 3.8.

**Lemma 3.10** (Middle application). *Let  $\rho, \sigma$  be arbitrary types.*

1. *Middle application is consistently defined for transitive right arguments, that is, if  $W \in \text{Con}_{\rho \rightarrow \sigma}$  and  $U \in \text{Con}_{\rho|L(W)}^{\text{ctr}}$ , then  $W \cdot U \in \text{Con}_\sigma$ .*
2. *Middle application is antimonotone in the right argument, in particular, if  $\Theta \in \text{Fin}_{\rho \rightarrow \sigma}$  and  $U, U' \in \text{Con}_\rho$  with  $U \vdash_\rho U'$ , then  $\Theta \cdot U \subseteq \Theta \cdot U'$ .*
3. *Middle application between neighborhoods is monotone in the left argument for transitive right arguments, that is, if  $W, W' \in \text{Con}_{\rho \rightarrow \sigma}$  are such that  $W \vdash_{\rho \rightarrow \sigma} W'$  and  $U \in \text{Con}_{\rho|L(W) \cup L(W')}^{\text{ctr}}$ , then  $W \cdot U \vdash_\sigma W' \cdot U$ .*

*Proof.* To show 1, let  $U \in \text{Con}_{\rho|L(W)}^{\text{ctr}}$  and  $b_1, b_2 \in W \cdot U$ . Then there exist  $U_1, U_2$  with  $\langle U_i, b_i \rangle \in W$  and  $U_i \asymp_\rho U$ . The transitivity of  $U$  implies  $U_1 \asymp_\rho U_2$ , and the consistency of  $W$  ensures that  $b_1 \asymp_\sigma b_2$ .

For 2, assume  $U$  and  $U'$  such that  $U \vdash_\rho U'$ , and let  $b \in \Theta \cdot U$ . There is some  $U_b$  such that  $\langle U_b, b \rangle \in \Theta$  and  $U_b \asymp_\rho U$ . By propagation (1) we get  $U_b \asymp_\rho U'$ , so  $b \in \Theta \cdot U'$ .

For 3. By 1, the assumption that  $U \in \text{Con}_{\rho|L(W) \cup L(W')}^{\text{ctr}}$  ensures that the result of both middle applications is a neighborhood (in general, if  $\Gamma \subseteq \Gamma'$  then  $\text{Con}_{\rho|\Gamma'}^{\text{ctr}} \subseteq \text{Con}_{\rho|\Gamma}^{\text{ctr}}$ ). Let  $b' \in W' \cdot U$ . By the definition of middle application, there exists a  $\langle U', b' \rangle \in W'$ , such that  $U' \asymp_\rho U$ . Since  $W \vdash_{\rho \rightarrow \sigma} W'$ , there is a subneighborhood  $\{\langle U_i, b_i \rangle \mid i = 1, \dots, m\} \subseteq W$ , such that for all  $i = 1, \dots, m$  we have  $U' \vdash_\rho U_i$  and  $\{b_i \mid i = 1, \dots, m\} \vdash_\sigma b'$ ; by propagation it follows that  $U \asymp_\rho U_i$  for all  $i$ . This means that  $b_i \in W \cdot U$  for all  $i$ , so  $W \cdot U \vdash_\sigma b'$ , and we are done.  $\square$

<sup>4</sup>The proof in fact shows that the entailment here is linear ( $U$  entails  $b$  linearly when  $\{a\} \vdash b$  for some  $a \in U$ , see [23]).

**Lemma 3.11.** *Let  $\Theta \in \text{Fin}_{\rho \rightarrow \sigma}$ .*

1. *For all  $U, U' \in \text{Con}_{L(\Theta)}^{\text{ctr}}$ , if  $U \asymp_{\rho} U'$  then  $\Theta \cdot U = \Theta \cdot U'$ .*
2. *For all  $U \in \text{Con}_{L(\Theta)}^{\text{ctr}}$ , we have  $\Theta \cdot U = \Theta \cdot \hat{U}$ .*
3. *For all  $U \in \text{Con}_{L(\Theta)}^{\text{max}}$  and  $U' \in \text{Con}_{\rho}$ , if  $U \asymp_{\rho} U'$  then  $\Theta \cdot U \subseteq \Theta \cdot U'$ .*

*Proof.* For the first statement, assume that  $U \asymp_{\rho} U'$  and let  $b \in R(\Theta)$ . We have  $b \in \Theta \cdot U$  if and only if there is some  $U_b$  with  $\langle U_b, b \rangle \in \Theta$  and  $U \asymp_{\rho} U_b$ . Since  $U$  is transitive in  $L(\Theta)$ , we get  $U' \asymp_{\rho} U_b$  by the assumption, so  $b \in \Theta \cdot U'$ . The converse is similar.

For the second statement, let  $U \in \text{Con}_{L(\Theta)}^{\text{ctr}}$ . By the definition of middle application, if  $b \in \Theta \cdot U$ , then there is some  $U_b$  with  $\langle U_b, b \rangle \in \Theta$ , such that  $U_b \asymp_{\rho} U$ ; since  $U$  is transitive, by the maximality of its maximal extension it follows that  $\hat{U} \vdash_{\rho} U_b$ , so the definition of application gives us  $b \in \Theta \cdot \hat{U}$ . For the other way around, if  $b \in \Theta \cdot \hat{U}$ , then there is a  $U_b$  with  $\langle U_b, b \rangle \in \Theta$ , such that  $\hat{U} \vdash_{\rho} U_b$ ; then  $U \asymp_{\rho} U_b$  by propagation (1), so  $b \in \Theta \cdot U$ , by the definition of middle application.

For the third statement, let  $U$  be maximal in  $L(\Theta)$  and  $U'$  some neighborhood with  $U \asymp_{\rho} U'$ . For every  $b \in \Theta \cdot U$ , by the definition of middle application, there is some  $U_b$  with  $\langle U_b, b \rangle \in \Theta$ , such that  $U \asymp_{\rho} U_b$ , which by maximality means that  $U \vdash_{\rho} U_b$ ; by propagation we get  $U' \asymp_{\rho} U_b$ , so  $b \in \Theta \cdot U'$ , and we are done.  $\square$

We close the section with a hint on how extremality evolves over types.

**Lemma 3.12.** *Let  $\Theta \in \text{Fin}_{\rho \rightarrow \sigma}$  and  $W \in \text{Con}_{\Theta}$ . We have  $W \in \text{Con}_{\Theta}^{\text{ctr}}$  if one of the following holds.*

1. *For all  $U \in \text{Con}_{L(\Theta)}^{\text{ctr}}$  we have  $W \cdot U \in \text{Con}_{\Theta \cdot U}^{\text{ctr}}$ .*
2. *For all  $U \in \text{Con}_{L(\Theta)}^{\text{max}}$  we have  $W \cdot U \in \text{Con}_{\Theta \cdot U}^{\text{ctr}}$ .*

*Proof.* For the first criterion, let  $\langle U_i, b_i \rangle \in \Theta$  be such that  $\langle U_i, b_i \rangle \asymp_{\rho \rightarrow \sigma} W$  for  $i = 1, 2$ , and assume that  $U_1 \asymp_{\rho} U_2$ . Consider a  $U \in \text{Con}_{L(\Theta)}^{\text{ctr}}$  with  $U \asymp_{\rho} U_1 \cup U_2$ ; then  $b_i \in \Theta \cdot U$  for each  $i$ . From  $\langle U_i, b_i \rangle \asymp_{\rho \rightarrow \sigma} W$ , by Lemma 3.10.1, we get  $b_i \asymp_{\sigma} W \cdot U$  for both  $i$ , so by the assumption we get  $b_1 \asymp_{\sigma} b_2$ .

To get the second criterion, it suffices to show that

$$\bigvee_{U \in \text{Con}_{L(\Theta)}^{\text{max}}} W \cdot U \in \text{Con}_{\Theta \cdot U}^{\text{ctr}} \rightarrow \bigvee_{U \in \text{Con}_{L(\Theta)}^{\text{ctr}}} W \cdot U \in \text{Con}_{\Theta \cdot U}^{\text{ctr}}.$$

Assume that  $W$  is such that  $W \cdot U \in \text{Con}_{\Theta \cdot U}^{\text{ctr}}$ , for all  $U \in \text{Con}_{L(\Theta)}^{\text{max}}$ , and let  $U \in \text{Con}_{L(\Theta)}^{\text{ctr}}$ . Consider the maximal extension  $\hat{U}$  of  $U$ . On the one hand we have  $\hat{U} \in \text{Con}_{L(\Theta)}^{\text{max}}$ , so  $W \cdot \hat{U} \in \text{Con}_{\Theta \cdot \hat{U}}^{\text{ctr}}$  by the assumption. On the other hand we have  $\hat{U} \in \text{Con}_{L(\Theta)}^{\text{ctr}}$  with  $\hat{U} \asymp_{\rho} U$ , so  $\Theta \cdot \hat{U} = \Theta \cdot U$  and  $W \cdot \hat{U} = W \cdot U$ , by Lemma 3.11.1. It follows that  $W \cdot U \in \text{Con}_{\Theta \cdot U}^{\text{ctr}}$ , so we can apply the previous criterion and we are done.  $\square$

## 4 Totality of neighborhoods

In this section we take the first two steps of the strategy that we outlined in the introduction. A total object of type  $\rho \rightarrow \sigma$  is represented by a possibly infinite token set that (a) is an ideal, that is, consistent and deductively closed, (b) admits all totals of type

$\rho$  as arguments—a property we think of as “omnicipation”, for lack of a less pompous synonym for *admission* (or *acceptance*) of *all*, that would be as grammatically smooth—, and (c) responds to every total argument with a total value at type  $\sigma$ . To bring the notion down to the finite level we dispose of half of the demand (a), namely, that the set of tokens be deductively closed, and we reinterpret “admittance” and “response” in (b) and (c) in terms of consistency rather than entailment. The first move, which is clearly dictated by the demand of finiteness, in some sense causes the reaction of the second move: what we lose by denying deductive closure we have to regain with the wider and more tolerant scope of consistency.

## 4.1 Finite density

At type  $\rho$ , a *side extension* of a neighborhood  $U$  will be any neighborhood  $U'$  which is consistent with  $U$ . We give a name to this rather mundane notion just to point to its intended use: trivially, if  $U'$  is a side extension of  $U$ , then  $U' \cup U$  is an extension of  $U$ , and this is exactly how we will work towards finding total extensions of neighborhoods.

**Lemma 4.1.** *Let  $\rho$  and  $\sigma$  be types. For every  $W \in \text{Con}_{\rho \rightarrow \sigma}$  and for every mapping  $V \mapsto V'$  of neighborhoods  $V \in \text{Con}_{\sigma}$  to side extensions thereof, the finite set  $\bigcup_{U \in \text{Con}_{L(W)}^{\text{ctr}}} \langle U, (W \cdot U)' \rangle$  is a side extension of  $W$ .*

*Proof.* To show the consistency of the finite set, let  $\langle U_i, b_i \rangle$  be such that  $U_i \in \text{Con}_{L(W)}^{\text{ctr}}$  and  $b_i \in (W \cdot U_i)'$ , for  $i = 1, 2$ . If  $U_1 \preceq_{\rho} U_2$ , then  $W \cdot U_1 = W \cdot U_2$  by Lemma 3.11.1, hence  $(W \cdot U_1)' = (W \cdot U_2)'$ , and  $b_1 \preceq_{\sigma} b_2$ .

To show the side extension, let  $\langle U, b \rangle \in W$  and  $\langle U', b' \rangle$  be such that  $U' \in \text{Con}_{L(W)}^{\text{ctr}}$  and  $b' \in (W \cdot U)'$ . If  $U \preceq_{\rho} U'$  then by the definition of middle application we have  $b \in W \cdot U$ . But  $W \cdot U \preceq_{\sigma} (W \cdot U)'$  by assumption, so  $b \preceq_{\sigma} b'$ .  $\square$

As we mentioned in section 1, a *total token* at a base type  $t$  is a token  $p \in \text{Tok}_t$  which consists exclusively of proper constructors; write  $p \in \text{Tok}_t^g$ . We have  $Cp_1 \cdots p_r \in \text{Tok}_t^g$  if and only if  $C$  is a proper constructor of arity  $r$  and  $p_i \in \text{Tok}_t^g$  for all  $i = 1, \dots, r$ . So  $\text{SB}^*0 \notin \text{Tok}_{\mathbb{D}}^g$  but  $\text{SB}10 \in \text{Tok}_{\mathbb{D}}^g$ . Define *total neighborhoods* inductively over types:

$$U \in \text{Con}_t^g := \exists_{p \in \text{Tok}_t^g} U \vdash_t p,$$

$$W \in \text{Con}_{\rho \rightarrow \sigma}^g := \forall_{P \in \text{Con}_{\rho}^g} W \cdot P \in \text{Con}_{\sigma}^g.$$

Examples of total neighborhoods are  $\{\text{B}0^*, \text{B}^*1\}$  and  $\{\text{B}0\text{S}0\}$  at type  $\mathbb{D}$ , and the neighborhood  $\{\langle\{0\}\rangle, 0, \langle\{\text{S}^*\}\rangle, \text{S}0\}$  at type  $\mathbb{N} \rightarrow \mathbb{N}$ , which we saw in the introduction. A type 2 example would be the neighborhood

$$X = \{\langle\langle\{\text{tt}\}\rangle, 0\rangle, \langle\langle\{\text{tt}\}\rangle, \text{S}^*\rangle, \text{S}0\}$$

at type  $(\mathbb{B} \rightarrow \mathbb{N}) \rightarrow \mathbb{N}$ : Let  $T$  be a total neighborhood at type  $\mathbb{B} \rightarrow \mathbb{N}$ ; for the total singleton  $\{\text{tt}\}$  of type  $\mathbb{B}$  we get  $T \cdot \{\text{tt}\} \vdash_{\mathbb{N}} \text{S}^m 0$  for some  $m \geq 0$ , so we either have  $T \cdot \{\text{tt}\} \preceq_{\mathbb{N}} 0$  or  $T \cdot \{\text{tt}\} \preceq_{\mathbb{N}} \text{S}^*$ , which by Lemma 3.7.2 means either  $T \preceq_{\mathbb{B} \rightarrow \mathbb{N}} \langle\{\text{tt}\}\rangle, 0$  or  $T \preceq_{\mathbb{B} \rightarrow \mathbb{N}} \langle\{\text{tt}\}\rangle, \text{S}^*$ ; in the first case we get  $X \cdot T = \{0\}$  and in the second case  $X \cdot T = \{\text{S}0\}$ , both of which are total neighborhoods at type  $\mathbb{N}$ , as we wanted.

The arguments of a higher-type total neighborhood form a finite set that can accept (in the sense of middle application) every total neighborhood of the source type, in a

way that ensures their safe (that is, consistency-respecting) allotment to an appropriate value. Define *weakly omniceptive finite sets* explicitly by

$$\Gamma \in \text{Fin}_\rho^{\text{wo}} := \bigvee_{P \in \text{Con}_\rho^{\text{g}}} \bigexists_{U^P \in \text{Con}_\Gamma^{\text{ctr}}} P \approx_\rho U^P.$$

At a type  $\rho \rightarrow \sigma$  call  $\Theta$  a (*strongly*) *omniceptive finite set*, and write  $\Theta \in \text{Fin}_{\rho \rightarrow \sigma}^{\text{o}}$ , if

$$\Theta \in \text{Fin}_{\rho \rightarrow \sigma}^{\text{wo}} \wedge \bigvee_{U \in \text{Con}_{L(\Theta)}^{\text{max}}} (U \in \text{Con}_\rho^{\text{g}} \wedge \Theta \cdot U \in \text{Fin}_\sigma^{\text{o}} \wedge \langle U, \Theta \cdot U \rangle \subseteq \Theta),$$

which intuitively says that, beyond weak omniception,  $\Theta$  must meet certain requirements of finite totality, preservation of omniception, and closure under middle application for each of its left maximals. By convention, we set  $\text{Fin}_i^{\text{o}} := \text{Fin}_i^{\text{wo}}$  for arbitrary base types. In the following we always use the term “omniceptive” (without a qualifier) to mean “strongly omniceptive” (which for base types is the same as “weakly omniceptive”). A further convention that we will often employ in subsequent arguments is to use lower case “g”, “wo”, and “o” as superscripts of terms: for example, we may write “ $U^{\text{g}}$ ” for a neighborhood that is to belong to  $\text{Con}^{\text{g}}$  and possibly relates to a previously discussed  $U$ , or “ $\Gamma^{\text{o}}$ ” for a finite set that is to belong to  $\text{Fin}^{\text{o}}$  and possibly relates to a previously introduced  $\Gamma$ .

Call a type  $\rho$  *finitely dense* if every neighborhood  $U$  at  $\rho$  has a total side extension  $U^{\text{g}}$ , and *finitely omniceptive* if every finite set  $\Gamma$  has an omniceptive extension  $\Gamma^{\text{o}}$ . Moreover, call it *finitely total-transitive* if every total neighborhood  $U$  is *transitive* (in  $\rho$ ). The latter just means that  $U_1 \approx_\rho U \approx_\rho U_2$  implies  $U_1 \approx_\rho U_2$  for all  $U_1, U_2 \in \text{Con}_\rho$ , and we write  $U \in \text{Con}_\rho^{\text{ctr}}$ . Here’s a lemma to set the intuition straight.

**Lemma 4.2** (Compactness of transitivity). *Let  $\rho$  be a type. A neighborhood is transitive in  $\rho$  if and only if it is transitive for every finite set of  $\rho$ .*

*Proof.* For the less trivial direction, let  $U \in \text{Con}_\rho$  be such that  $U \in \text{Con}_{\rho|\Gamma}^{\text{ctr}}$  for every  $\Gamma \in \text{Fin}_\rho$ , and let  $U_1, U_2 \in \text{Con}_\rho$  be such that  $U_1 \approx_\rho U \approx_\rho U_2$ . Set  $\Gamma := U_1 \cup U \cup U_2$ ; then we have  $U_1 \approx_\rho U_2$  by  $\Gamma$ -transitivity.  $\square$

To start off the main argument we need some elementary definitions and observations concerning base types. The *size*  $\|a\|$  of a base-type token  $a \in \text{Tok}_i$  counts the proper constructors of the token:  $\|*\| = 0$  and  $\|Ca_1 \cdots a_r\| = 1 + \|a_1\| + \cdots + \|a_r\|$ . It is an easy induction to show that  $\{a\} \vdash_i b$  implies  $\|a\| \geq \|b\|$  for all tokens  $a, b$ . The *supremum* or *eigentoken*  $\text{sup}(U)$  of a base-type neighborhood  $U \in \text{Con}_i$  is defined by  $\text{sup}(\emptyset_i) = *_i$ ,  $\text{sup}(\{a\}) = a$ , and  $\text{sup}(\{a_1, \dots, a_m\}) = \text{sup}^t(\cdots \text{sup}^t(a_1, a_2) \cdots, a_m)$  for  $m > 1$ , where  $\text{sup}^t(a, *) = a$  and  $\text{sup}^t(Ca_1 \cdots a_r, Cb_1 \cdots b_r) = C\text{sup}^t(a_1, b_1) \cdots \text{sup}^t(a_r, b_r)$  (we do not need to define the auxiliary mapping  $\text{sup}^t$  on inconsistent pairs). Again, an easy induction shows that  $U \sim_i \{\text{sup}(U)\}$  for all  $U \in \text{Con}_i$ ; in particular, we can represent every total neighborhood  $P \in \text{Con}_i^{\text{g}}$  by its total eigentoken  $\text{sup}(P) \in \text{Tok}_i^{\text{g}}$ .

**Proposition 4.3.** *Every base type is finitely total-transitive, finitely dense, and finitely omniceptive.*

*Proof.* Let  $i$  be any base type with a distinguished nullary constructor  $0$ . For the transitivity of total neighborhoods, let  $P \in \text{Con}_i^{\text{g}}$  and  $U_1, U_2 \in \text{Con}_i$  be such that  $U_i \approx_i P$  for each  $i$ . Then  $P \vdash_i U_i$ , for both  $i = 1, 2$ , since, as is easy to see, total tokens are maximal at base types, so  $U_1 \approx_i U_2$ .

We turn to finite density by firstly considering tokens: the trivial token  $*$  is consistent with  $0$ , and if  $p_1, \dots, p_r$  are total tokens consistent with  $a_1, \dots, a_r$  respectively, then  $Cp_1 \cdots p_r$  is a total token consistent with  $Ca_1 \cdots a_r$ , for an  $r$ -ary constructor  $C$ . Finally, if  $U \in \text{Con}_t$ , and  $p$  is a total token consistent to  $\text{sup}(U)$ , then the neighborhood  $\{p\}$  is obviously a total neighborhood consistent with (above, even)  $U$ .

Before we turn to the finite omniception we need some auxiliary facts. Firstly, we claim that

$$\forall_{a \in \text{Tok}_t} \forall_{p \in \text{Tok}_t^g} (\|a\| > \|p\| \rightarrow a \not\asymp_t p). \quad (3)$$

Indeed, let  $p$  be a total token and  $a$  an arbitrary token. In case  $p$  consists of a single nullary constructor, then  $\|p\| = 1$  and there have to exist a constructor  $C$  with arity  $r > 0$  and further tokens  $a_1, \dots, a_r$  such that  $a = Ca_1 \cdots a_r$  (with  $\|a_i\| > 0$  for at least some  $i$ ); we have  $a \not\asymp_t p$  by the definition of consistency. In case  $p = Cp_1 \cdots p_r$  for some total tokens  $p_1, \dots, p_r$ , then  $a$  will either start with a different (non-nullary) head constructor than  $C$ , in which case we are done, or there must exist  $a_1, \dots, a_r$  such that  $a = Ca_1 \cdots a_r$ ; from  $\|a\| > \|p\|$  it follows that  $\sum_i \|a_i\| > \sum_i \|p_i\|$ , so there must exist at least some  $i$  with  $\|a_i\| > \|p_i\|$ ; by the induction hypothesis we know that this means that  $a_i \not\asymp_t p_i$ , therefore  $a \not\asymp_t p$ .

Now, for a fixed total token  $p \in \text{Tok}_t^g$  and a fixed natural number  $n$ , let  $U^{p,n}$  stand for the finite set  $\{a \in \text{Tok}_t \mid a \asymp_t p \wedge \|a\| \leq n\}$ ; this is a neighborhood thanks to the transitivity of the total  $p$ . Note that for every  $p$  we have  $U^{p,0} = \{*_t\}$ , and also, due to (3), for all  $n \geq \|p\|$  we have  $U^{p,n} = U^{p,\|p\|}$ . Now, either  $p$  consists of a single nullary constructor or not; in case it does, then  $U^{p,1} = \{*_t, p\}$ , while if  $p = Cp_1 \cdots p_r$  for a constructor  $C$  with arity  $r > 0$  and total tokens  $p_1, \dots, p_r$ , then we claim that the component neighborhoods satisfy

$$\bigvee_{i=1}^r U^{p,n}(i) = U^{p_i, n-1}. \quad (4)$$

Indeed, fix an  $i$ . From left to right, assume that  $a^i \in U^{p,n}(i)$ ; this means that there exists an  $a \in U^{p,n}$  such that  $a(i) = a^i$ ; by the definition of  $U^{p,n}$ , we know that  $a \asymp_t p$  which implies that  $a^i \asymp_t p_i$ ; we also know that  $\|a\| \leq n$ , so  $\|a^i\| \leq \|a\| - 1 \leq n - 1$ , therefore  $a^i \in U^{p_i, n-1}$ . For the other way around, assume that  $a^i \in U^{p_i, n-1}$ , that is, that  $a^i \asymp_t p_i$  and  $\|a^i\| \leq n - 1$ ; consider the token  $a := C \overrightarrow{*} a^i \overrightarrow{*}$  (which has  $a^i$  as its  $i$ -th component token and stars everywhere else); it obviously satisfies  $a \asymp_t p$ , and since  $\|a^i\| \leq n - 1$ , it also satisfies  $\|a\| \leq n$ , therefore  $a \in U^{p,n}$ ; it follows that  $a(i) = a^i \in U^{p,n}(i)$ .

Finally, we claim that the following holds for a fixed  $n$ :

$$\forall_{a \in \text{Tok}_t} \forall_{p \in \text{Tok}_t^g} (\|a\| = n \leq \|p\| \wedge a \asymp_t U^{p,n} \rightarrow U^{p,n} \vdash_t a). \quad (5)$$

Indeed, let  $p$  be some total token and  $a$  an arbitrary one, such that  $\|a\| = n \leq \|p\|$  and  $a \asymp_t U^{p,n}$ . In case  $n = 0$ , we necessarily have  $a = *_t \in U^{p,0}$ . In case  $n > 0$ , then for some constructor  $C$  of the algebra  $t$  with arity  $r$  there exist total tokens  $p_1, \dots, p_r$  such that  $p = Cp_1 \cdots p_r$ , and since  $C \overrightarrow{*} \in U^{p,n}$  and  $a \asymp_t U^{p,n}$ , there exist tokens  $a_1, \dots, a_r$  such that  $a = Ca_1 \cdots a_r$ ; for every  $i = 1, \dots, r$ , since  $a \asymp_t U^{p,n}$  we have  $a_i \asymp_t U^{p,n}(i)$ , which by (4) means that  $a_i \asymp_t U^{p_i, n-1}(i)$ , while from the assumption that  $\|a\| = n$  we get  $\|a_i\| \leq n - 1$ ; the induction hypothesis yields  $U^{p_i, n-1} \vdash_t a_i$  for every  $i$ ; so we get

$$U^{p,n} \sim_t CU^{p,n}(1) \cdots U^{p,n}(r) \stackrel{(4)}{=} CU^{p_1, n-1} \cdots U^{p_r, n-1} \vdash_t Ca_1 \cdots a_r = a,$$

and we are done.

Now for the finite omniception. If  $\Gamma$  is trivial (that is, if it carries no proper information), then set  $\Gamma^o := \{*\}$ . If not, let  $m := \max\{\|\text{sup}(U)\| \mid U \in \text{Con}_\Gamma\}$ , and set  $\Gamma^o := \{a \in \text{Tok}_t \mid \|a\| \leq m\}$ . Consider a total neighborhood  $P$  and let  $p := \text{sup}(P)$ . In case  $\|p\| \leq m$  we have  $p \in \Gamma^o$ , so it suffices to set  $U^P := \{p\}$ . In case  $\|p\| > m$ , we set  $U^P := U^{p,m}$ ; by the construction of  $\Gamma^o$  we have  $U^{p,m} \in \text{Con}_{\Gamma^o}$ , and we also have that  $U^{p,m}$  is transitive in  $\Gamma^o$ : for any two tokens  $b_1, b_2 \in \Gamma^o$  with  $b_1 \asymp_1 U^{p,m} \asymp_1 b_2$ , since  $\|b_i\| \leq m < \|p\|$  for both  $i$ , we get  $b_i \in U^{p,m} \vdash_1 b_i$  from (5), so  $b_1 \asymp_1 b_2$ .  $\square$

**Proposition 4.4** (Finite total-transitivity). *Let  $\rho$  and  $\sigma$  be finitely total-transitive types. If  $\rho$  is finitely dense then  $\rho \rightarrow \sigma$  is finitely total-transitive.*

*Proof.* Let  $T \in \text{Con}_{\rho \rightarrow \sigma}^g$  and  $W_1, W_2 \in \text{Con}_{\rho \rightarrow \sigma}$ , with  $W_1 \asymp_{\rho \rightarrow \sigma} T \asymp_{\rho \rightarrow \sigma} W_2$ . Consider pairs  $\langle U_i, b_i \rangle \in W_i$ ,  $i = 1, 2$ , and assume that  $U_1 \asymp_\rho U_2$ . By the finite density at  $\rho$ , there exists a  $P \in \text{Con}_\rho^g$ , such that  $P \asymp_\rho U_1 \cup U_2$ . By the assumptions at  $\rho$  and Lemma 3.10.1 we get  $b_1 \asymp_\sigma T \cdot P \asymp_\sigma b_2$ . But  $T \cdot P$  is total, so the assumption at  $\sigma$  gives  $b_1 \asymp_\sigma b_2$ .  $\square$

**Proposition 4.5** (Finite density). *Let  $\rho$  and  $\sigma$  be types. If  $\rho$  is finitely omniceptive and  $\sigma$  finitely dense and finitely total-transitive, then  $\rho \rightarrow \sigma$  is finitely dense.*

*Proof.* Let  $W \in \text{Con}_{\rho \rightarrow \sigma}$  be any neighborhood. By finite omniception at  $\rho$  we get a  $\Gamma \in \text{Fin}_\rho^o$  with  $L(W) \subseteq \Gamma$ . Consider the neighborhood  $W^o := \bigcup_{U \in \text{Con}_\Gamma} \langle U, W \cdot U \rangle$ ; by Lemma 3.9 we have  $W \sim_{\rho \rightarrow \sigma} W^o$ . Now set

$$W^g := \bigcup_{U \in \text{Con}_{L(W^o)}^{\text{ctr}}} \langle U, (W^o \cdot U)^g \rangle,$$

with the help of density at  $\sigma$ ; note that  $L(W^g) = L(W^o) = \Gamma$ . This is a side extension of  $W^o$  (therefore of  $W$  as well) by Lemma 4.1.

To show that it is total, let  $P \in \text{Con}_\rho^g$ . Since  $L(W^g)$  is omniceptive (in fact, that it is weakly omniceptive is enough), there is some  $U^P \in \text{Con}_{L(W^g)}^{\text{ctr}}$  such that  $P \asymp_\rho U^P$ . We have  $\langle U^P, (W^o \cdot U^P)^g \rangle \subseteq W^g$  by construction, and  $W^g \cdot P = (W^o \cdot U^P)^g$ , since, by transitivity of total neighborhoods at  $\sigma$ , the value  $W^o \cdot P$  is independent from the choice of  $U^P$ .  $\square$

**Proposition 4.6** (Finite omniception). *Let  $\rho$  and  $\sigma$  be finitely total-transitive types. If  $\rho$  is finitely dense and  $\sigma$  finitely omniceptive, then  $\rho \rightarrow \sigma$  is finitely omniceptive.*

*Proof.* Let  $\Theta \in \text{Fin}_{\rho \rightarrow \sigma}$  be any finite set. Extend it as follows:

$$\Theta^o := \Theta \cup \bigcup_{U \in \text{Con}_{L(\Theta)}^{\text{max}}} \langle U^g, (\Theta \cdot U)^o \rangle,$$

with the use of finite density at  $\rho$  and finite omniception at  $\sigma$ .

If we show that this is weakly omniceptive, then it will be omniceptive immediately by construction (based on Lemma 3.5). Let  $T \in \text{Con}_{\rho \rightarrow \sigma}^g$ . For every  $U \in \text{Con}_{L(\Theta^o)}^{\text{max}}$  we have  $T \cdot U^g \in \text{Con}_\sigma^g$ , and since  $(\Theta \cdot U)^o$  is omniceptive, there will be some  $V^{T \cdot U^g} \in \text{Con}_{(\Theta \cdot U)^o}^{\text{ctr}}$ , such that  $T \cdot U^g \asymp_\sigma V^{T \cdot U^g}$  ( $\star$ ). Fix these side extensions and set

$$W^T := \bigcup_{U \in \text{Con}_{L(\Theta^o)}^{\text{max}}} \langle U^g, V^{T \cdot U^g} \rangle.$$

We have  $W^T \subseteq \Theta^o$  by construction. Moreover, we have  $T \asymp_{\rho \rightarrow \sigma} W^T$ : let  $\langle U, b \rangle \in T$  and  $\langle U', b' \rangle \in W^T$  be such that  $U \asymp_\rho U'$ ; we have  $b \in T \cdot U'$  and  $b' \in V^{T \cdot U'}$ , so  $b \asymp_\sigma b'$  by

( $\star$ ). Since  $T$  is total,  $W^T$  is a neighborhood by transitivity of total neighborhoods, which we get for  $\rho \rightarrow \sigma$  by Proposition 4.4. Finally, it is transitive in  $\Theta^o$  by Lemma 3.12.2, since for every  $U \in \text{Con}_{L(\Theta)}^{\max}$  we have by construction  $W^T \cdot U = V^{T \cdot U}$ , which is transitive in  $\Theta \cdot U$  by omniception.  $\square$

**Theorem 4.7.** *Every type is finitely omniceptive, finitely total-transitive, and, in particular, finitely dense.*

*Proof.* We get this by mutual induction over types from Propositions 4.3, 4.4, 4.5, and 4.6.  $\square$

## 4.2 Totality of transitive neighborhoods

There is plenty of evidence to suggest that total neighborhoods at  $\rho$  are to  $\text{Con}_\rho$  what transitive neighborhoods in  $\Gamma$  are to  $\text{Con}_\Gamma$ . For one, Theorem 4.7 shows that total neighborhoods are transitive. Furthermore, an immediate corollary of total transitivity is that consistency, restricted to the total neighborhoods, becomes an equivalence relation, that is,

$$\bigvee_{P_1, P_2, P_3 \in \text{Con}_\rho^g} (P_1 \asymp_\rho P_2 \asymp_\rho P_3 \rightarrow P_1 \asymp_\rho P_3) .$$

Here are further examples of using total transitivity, which include some more evidence to this effect.

**Lemma 4.8.** *Let  $\rho$  and  $\sigma$  be types. Let  $\Theta \in \text{Fin}_{\rho \rightarrow \sigma}$ ,  $P, P' \in \text{Con}_\rho^g$ , and  $U, U' \in \text{Con}_\rho$ .*

1. *For every  $U^P \in \text{Con}_{L(\Theta)}$  with  $P \asymp_\rho U^P$ , we have  $\Theta \cdot P \subseteq \Theta \cdot U^P$ . Moreover, we have  $\Theta \cdot P = \Theta \cdot U^P$  whenever  $U^P \in \text{Con}_{L(\Theta)}^{\max}$ .*
2. *If  $P \asymp_\rho P'$  then  $\Theta \cdot P = \Theta \cdot P'$ .*
3. *If  $U \in \text{Con}_\rho^g$  and  $U' \vdash_\rho U$ , then  $U' \in \text{Con}_\rho^g$ .*
4. *If  $T \in \text{Con}_{\rho \rightarrow \sigma}^g$  and  $U \in \text{Con}_{L(T)}^{\text{tr}}$  then  $T \cdot U \in \text{Con}_\sigma^g$ .*

*Proof.* For 1, let  $b \in \Theta \cdot P$ . Then, by the definition of middle application, there is some  $U$  with  $\langle U, b \rangle \in \Theta$ , such that  $U \asymp_\rho P$ . From  $U \asymp_\rho P \asymp_\rho U^P$  we get  $U \asymp_\rho U^P$  by Theorem 4.7, so the definition of middle application yields that  $b \in \Theta \cdot U^P$ . Moreover, if  $U^P$  is actually maximal in  $L(\Theta)$ , then by Lemma 3.11.3 we immediately get  $\Theta \cdot U^P \subseteq \Theta \cdot P$ .

For 2, assume that  $P \asymp_\rho P'$  and let  $b \in \Theta \cdot P$ . By the definition of middle application, there is some  $U$  with  $\langle U, b \rangle \in \Theta$ , such that  $U \asymp_\rho P$ . By Theorem 4.7 and the assumption we have  $U \asymp_\rho P'$ , so  $b \in \Theta \cdot P'$ .

For 3. At a base type  $\iota$  if  $U \in \text{Con}_\iota^g$ , then there exists a total token  $p$  such that  $U \vdash_\iota p$ . The transitivity of entailment yields what we need. At a higher type  $\rho \rightarrow \sigma$ , let  $W \in \text{Con}_{\rho \rightarrow \sigma}^g$  and  $W' \vdash_{\rho \rightarrow \sigma} W$ . Let further  $P \in \text{Con}_\rho^g$ . By Theorem 4.7,  $P$  is transitive for  $L(W)$ , so by the left monotonicity of middle application on transitive arguments (Lemma 3.10.3), we have  $W' \cdot P \vdash_\sigma W \cdot P$ , and by the totality of  $W$  we get  $W \cdot P \in \text{Con}_\sigma^g$ , so the induction hypothesis at  $\sigma$  finishes the job.

For 4. By Theorem 4.7 there exists some  $P^U \in \text{Con}_\rho^g$  with  $P^U \asymp_\rho U$ . By 1 we have  $T \cdot P^U \subseteq T \cdot U$ , where  $T \cdot U$  is consistent by Lemma 3.10.1. It follows by 3 that  $T \cdot U \in \text{Con}_\sigma^g$ .  $\square$

Note in particular that Lemma 4.8.3 is analogous to Lemma 3.3 (both of them actually anticipate Lemma 5.2).

We now show that the correspondence between transitivity and finite totality is complete.

**Theorem 4.9** (Explicit finite totality). *At every type, a neighborhood is total if and only if it is transitive.*

*Proof by induction over types.* The rightward direction we have of course from Theorem 4.7. For the other direction, we have to show that, at each type, every transitive neighborhood must be total.

At a base type  $\iota$ , assume that  $U \in \text{Con}_\iota^{\text{ctr}}$ . Obviously, we have  $U \not\sim_\iota \{*\}$ , so there will be a constructor  $C$  and tokens  $a_1, \dots, a_r \in \text{Tok}_\iota$  such that  $U \sim_\iota \{Ca_1 \cdots a_r\}$  (its supremum). By Lemma 3.4.2, since  $U$  is transitive, for any two tokens  $b_1, b_2 \in \text{Tok}_\iota$  we will have  $b_1 \succ_\iota U \succ_\iota b_2$  imply  $b_1 \succ_\iota b_2$ . Then for  $i = 1, \dots, r$  we have

$$\begin{aligned} b_{1i} \succ_\iota a_i \succ_\iota b_{2i} &\Rightarrow Ca_1 \cdots b_{1i} \cdots a_r \succ_\iota U \succ_\iota Ca_1 \cdots b_{2i} \cdots a_r \\ &\stackrel{\text{ctr}}{\Rightarrow} Ca_1 \cdots b_{1i} \cdots a_r \succ_\iota Ca_1 \cdots b_{2i} \cdots a_r \\ &\Rightarrow b_{1i} \succ_\iota b_{2i} \end{aligned}$$

for any  $b_{1i}, b_{2i} \in \text{Tok}_\iota$ , which by induction hypothesis yields  $a_i \in \text{Tok}_\iota^g$ . It follows that  $Ca_1 \cdots a_r$  itself is a total token, so  $U$  is a total neighborhood.

At type  $\rho \rightarrow \sigma$ , assume that  $W \in \text{Con}_{\rho \rightarrow \sigma}^{\text{ctr}}$ , and let  $P \in \text{Con}_\rho^g$ . For any  $b_1, b_2 \in \text{Tok}_\sigma$  we have

$$\begin{aligned} b_1 \succ_\sigma W \cdot P \succ_\sigma b_2 &\Leftrightarrow \langle P, b_1 \rangle \succ_{\rho \rightarrow \sigma} W \succ_{\rho \rightarrow \sigma} \langle P, b_2 \rangle \\ &\stackrel{\text{ctr}}{\Rightarrow} \langle P, b_1 \rangle \succ_{\rho \rightarrow \sigma} \langle P, b_2 \rangle \\ &\Rightarrow b_1 \succ_\sigma b_2, \end{aligned}$$

which means that  $W \cdot P$  is transitive in  $\sigma$ , so by the induction hypothesis at  $\sigma$  we get  $W \cdot P \in \text{Con}_\sigma^g$ , and by the definition of finite totality we have  $W \in \text{Con}_{\rho \rightarrow \sigma}^g$ , as we wanted.  $\square$

The theorem indicates that our notion of finite totality is a robust one. In the next section we will see how we can connect it to the traditional notion of totality for ideals. Interestingly, we will see that its equivalence to transitivity is peculiar to the finitary level: in Proposition 5.12 the respective correspondence for ideals is shown to be tilted.

## 5 Elevating totality to ideals

The last step in our strategy is to find a canonical extension of a total neighborhood to a total ideal. The natural candidate would be the deductive closure of a neighborhood, but again, closure under entailment is too strict for our purposes, since it presents recurring technical difficulties stemming from the fact that the application of ideals is an ‘‘upper’’ one, while we have defined finite totality in terms of middle application. Instead, based on the transitivity of total neighborhoods, using *closure under consistency* proves to be a more natural choice.

## 5.1 Density

The notion of continuity that we employ in our setting implies that if we are given an estimate  $V$  on a value  $f(x)$  then we can find an adequate estimate  $U_V$  on the argument  $x$  of  $f$ ; let us highlight this elementary fact since we will need it later on.

**Lemma 5.1** (Finite support). *Let  $f : \rho \rightarrow \sigma$  and  $x : \rho$ . For every  $V \in \text{Con}_\sigma$  with  $V \subseteq f(x)$  there exists a  $U_V \in \text{Con}_\rho$  such that  $U_V \subseteq x$  and  $\langle U_V, V \rangle \subseteq f$ .*

*Proof.* From (FS) it follows directly that if  $b \in f(x)$  then there exists a  $U_b \subseteq x$  such that  $\langle U_b, b \rangle \in f$  due to the deductive closure of  $f$ . Assuming then that  $V$  is such that  $V \subseteq f(x)$ , for  $U_V := \bigcup_{b \in V} U_b$  we indeed have  $U_V \subseteq x$ , and also  $\langle U_V, V \rangle \subseteq f$  by the deductive closure of  $f$ .  $\square$

An ideal  $x : \rho$  is a *total ideal*, for which we write  $G_\rho(x)$  or  $x \in G_\rho$ , if it conforms to the following inductive clauses.

$$G_\iota(x) := \exists_{P \in \text{Con}_\iota^g} P \subseteq x,$$

$$G_{\rho \rightarrow \sigma}(f) := \forall_{x:\rho} (G_\rho(x) \rightarrow G_\sigma(fx)).$$

Note that the base type definition is equivalent to demanding the existence of a  $p \in \text{Tok}_\iota^g$  such that  $p \in x$ . Totality of ideals is upwards closed.

**Lemma 5.2** (Extension lemma). *At type  $\rho$ , if  $x, y : \rho$  are such that  $G_\rho(x)$  and  $x \subseteq y$  then  $G_\rho(y)$ .*

*Proof.* At a base type  $\iota$ , let  $G_\iota(x)$  and  $y : \iota$  be two ideals with  $x \subseteq y$ . Then there is a total token  $p \in \text{Tok}_\iota^g$ , such that  $p \in x$ , so also  $p \in y$ . At a higher type  $\rho \rightarrow \sigma$ , let  $G_{\rho \rightarrow \sigma}(f)$ ,  $g : \rho \rightarrow \sigma$ , and assume that  $f \subseteq g$ . We want to show that  $g$  is also total, so consider an arbitrary  $x$  with  $G_\rho(x)$ . By the totality of  $f$  we have that  $G_\sigma(fx)$ , and since it is straightforward to see that  $fx \subseteq gx$ , we get  $G_\sigma(gx)$  by the induction hypothesis at  $\sigma$ .  $\square$

The main argument starts with the following obvious observation.

**Lemma 5.3.** *At every type, if a neighborhood is transitive then its consistency closure is an ideal (and the converse holds as well).*

*Proof.* Let  $\rho$  be a type and  $U \in \text{Con}_\rho^{\text{ctr}}$ . By transitivity, every two tokens in the consistency closure of  $U$  will be consistent, and the consistency closure is already deductively closed:  $U' \subseteq \tilde{U}$  means  $U \asymp_\rho U'$  by definition, so if  $U' \vdash_\rho a$ , then propagation yields  $U \asymp_\rho a$ , hence  $a \in \tilde{U}$  as well. The converse is also direct to show.  $\square$

By Theorem 4.9, an immediate consequence of this lemma is that the consistency closure of a total neighborhood is an ideal, so it suffices to show that, for a given  $P \in \text{Con}_\rho^g$ , we

must have  $G_\rho(\tilde{P})$ . Consider the following statements for an arbitrary type  $\rho$ .

$$\forall \Gamma \in \text{Fin}_\rho^o \quad \forall x \in G_\rho \quad \exists U^x \in \text{Con}_\rho^{\text{ctr}} \quad U^x \asymp_\rho x, \quad (\text{O})$$

$$\forall U, U' \in \text{Con}_\rho \quad \forall x \in G_\rho \quad (U' \not\asymp_\rho U \asymp_\rho x \rightarrow \exists U_0 \in \text{Con}_\rho \quad (U \vdash_\rho U_0 \subseteq x \wedge U' \not\asymp_\rho U_0)), \quad (\text{W})$$

$$\forall P \in \text{Con}_\rho^g \quad \tilde{P} \in G_\rho, \quad (\text{C})$$

$$\forall U \in \text{Con}_\rho \quad \exists x \in G_\rho \quad U \subseteq x. \quad (\text{D})$$

The first one is an expression of infinitary *omniception*, as it states that an omniceptive finite set accepts each total *ideal* by being consistent with it with one of its transitive neighborhoods. The second expresses *inconsistency preserving witnessing* of the consistency between a total ideal and a neighborhood; the claimed witness is stronger than the neighborhood itself, since it lies below both the total ideal and the neighborhood, and in a sense to be made clearer after Lemma 5.8 below, it provides the missing feature from omniception that we need to achieve totality on the level of ideals. The third one is the crux of our strategy, as it says that the *consistency closure* of a total neighborhood is a total ideal, and the fourth one, of course, is *density*.

**Proposition 5.4** (Conditional density). *Let  $\rho$  be a type. If (C) holds in  $\rho$  then also (D) holds in  $\rho$ .*

*Proof.* Let  $U$  be any neighborhood at type  $\rho$ . By Theorem 4.7 there exists a total neighborhood  $P^U$  such that  $U \asymp_\rho P^U$ . Then  $U \subseteq \tilde{P}^U$  by definition, whereas  $\tilde{P}^U \in G_\rho$  by (C). We set  $x := \tilde{P}^U$  and we are done.  $\square$

**Lemma 5.5.** *Every base type satisfies (O), (W), (C), and (D).*

*Proof.* Let  $\iota$  be some base type. To show (O), consider an omniceptive finite set  $\Gamma$  and a total ideal  $x$ . By the totality of  $x$  there is some  $P \in \text{Con}_\iota^g$  such that  $P \subseteq x$ , and by the omniception of  $\Gamma$  there is some  $U^P \in \text{Con}_\Gamma^{\text{ctr}}$  such that  $U^P \asymp_\iota P$ . Set  $U^x := U^P$ . Then for every  $U \subseteq x$  we have  $U^x \asymp_\iota P \asymp_\iota U$ , which implies  $U^x \asymp_\iota U$  by the total transitivity of  $\iota$  (Proposition 4.3), so  $U^x \asymp_\iota x$ .<sup>5</sup>

To show (W), let  $U$  and  $U'$  be neighborhoods and  $x$  be a total ideal, such that  $U' \not\asymp_\iota U \asymp_\iota x$ . By the totality of  $x$  there exists a total neighborhood  $P$  such that  $P \subseteq x$ . We have of course  $P \asymp_\iota U$ , which, since total tokens are maximal at base types, implies that  $P \vdash_\iota U$ . This in turn implies that  $U \subseteq x$  by the deductive closure of  $x$ , so we may set  $U_0 := U$ , which trivially meets the stated requirements.

To show (C), let  $P$  be some total neighborhood. Then there is some total token  $p \in \text{Tok}_\iota$  with  $P \vdash p$ ; a fortiori we have  $P \asymp_\iota p$ , so  $p \in \tilde{P}$  by the definition of consistency closure. Since by Lemma 5.3 the set  $\tilde{P}$  is an ideal, we conclude that it is in fact total.

Finally, that every base type is dense we get from Proposition 5.4, since (C) already holds.  $\square$

**Proposition 5.6** (Omniception). *Let  $\rho$  and  $\sigma$  be types. If (C) holds in  $\rho$  and (O) holds in  $\sigma$  then (O) holds in  $\rho \rightarrow \sigma$ .*

<sup>5</sup>Notice that here we only needed *weak* omniception from  $\Gamma$ . Furthermore, observe that in the flat setting this argument would fail due to the requirement of finiteness of  $\Gamma$ .

*Proof.* Let  $\Theta \in \text{Fin}_\rho^o \rightarrow \sigma$  and  $f \in G_{\rho \rightarrow \sigma}$ . By the finite omniception of  $\Theta$  we know that each  $U \in \text{Con}_{L(\Theta)}^{\max}$  is a total neighborhood, so by (C) at  $\rho$  we have  $\tilde{U} \in G_\rho$ . By the totality of  $f$  we have that  $f(\tilde{U}) \in G_\sigma$ , so there will be some  $V^{f(\tilde{U})} \in \text{Con}_{\Theta \cdot U}^{\text{ctr}}$  such that  $V^{f(\tilde{U})} \asymp_\sigma f(\tilde{U})$ , because  $\Theta \cdot U$  is omniceptive by the finite omniception of  $\Theta$  and (O) at  $\sigma$ . Based on these, we may set

$$W^f := \bigcup_{U \in \text{Con}_{L(\Theta)}^{\max}} \langle U, V^{f(\tilde{U})} \rangle.$$

We have  $W^f \in \text{Con}_\Theta^{\text{ctr}}$  by Lemma 3.12.2. Furthermore, let  $\langle U_0, b_0 \rangle \in W$  and  $\langle U, b \rangle \in f$  be such that  $U \asymp_\rho U_0$ ; then  $U \subseteq \tilde{U}_0$  (remember that  $U_0$  is a total neighborhood) and consequently  $\bar{U} \subseteq \tilde{U}_0$  by the propagation of consistency; by the monotonicity of  $f$  we get  $f(\bar{U}) \subseteq f(\tilde{U}_0)$ , so since  $b \in f(\bar{U})$  it must also be  $b \in f(\tilde{U}_0)$ ; but  $f(\tilde{U}_0) \asymp_\sigma V^{f(\tilde{U}_0)}$  and  $b_0 \in V^{f(\tilde{U}_0)}$ , so  $b \asymp_\sigma b_0$ , as we wanted.  $\square$

**Proposition 5.7 (Witnessing).** *Let  $\rho$  and  $\sigma$  be types. If (D) holds in  $\rho$  and (W) holds in  $\sigma$  then (W) holds in  $\rho \rightarrow \sigma$ .*

*Proof.* Let  $f \in G_{\rho \rightarrow \sigma}$  and  $W, W' \in \text{Con}_{\rho \rightarrow \sigma}$  be such that  $W' \not\asymp_{\rho \rightarrow \sigma} W \asymp_{\rho \rightarrow \sigma} f$ . For  $i = 1, \dots, m$ , let  $U'_i \in \text{Con}_{L(W')}$  and  $U_i \in \text{Con}_{L(W)}$  run through all witnessing pairs of inconsistency between  $W'$  and  $W$ , that is, cover all the cases where

$$U'_i \asymp_\rho U_i \wedge W' \cdot U'_i \not\asymp_\sigma W \cdot U_i.$$

By (D) at  $\rho$ , for each  $i$  there exists an  $x_i \in G_\rho$  such that  $U'_i \cup U_i \subseteq x_i$ . By the consistency of (upper) application, for every such  $x_i$  we have  $W \cdot U_i \asymp_\sigma f(x_i)$ , and by (W) at  $\sigma$  there exists some  $V_{i0} \in \text{Con}_\sigma$  such that  $W \cdot U_i \vdash_\sigma V_{i0} \subseteq f(x_i)$  and  $W' \cdot U'_i \not\asymp_\sigma V_{i0}$ . By Lemma 5.1, there exists some  $U_{V_{i0}} \subseteq x_i$  for every  $i$  such that  $\langle U_{V_{i0}}, V_{i0} \rangle \subseteq f$ . Letting  $U_{i0} := U_{V_{i0}} \cup U'_i \cup U_i$ , by the deductive closure of  $f$  it follows that  $\langle U_{i0}, V_{i0} \rangle \subseteq f$ . Since by the hypotheses at  $\sigma$  for every  $i$  we have

$$\langle U_i, W \cdot U_i \rangle \vdash_{\rho \rightarrow \sigma} \langle U_{i0}, V_{i0} \rangle \subseteq f \wedge \langle U'_i, W' \cdot U'_i \rangle \not\asymp_{\rho \rightarrow \sigma} \langle U_{i0}, V_{i0} \rangle,$$

it follows that

$$W \vdash_{\rho \rightarrow \sigma} \bigcup_{i=1}^m \langle U_{i0}, V_{i0} \rangle \subseteq f \wedge W' \not\asymp_{\rho \rightarrow \sigma} \bigcup_{i=1}^m \langle U_{i0}, V_{i0} \rangle,$$

so we may set  $W_0 := \bigcup_{i=1}^m \langle U_{i0}, V_{i0} \rangle$  and we are done.  $\square$

We may generalize the property (W) to account for inconsistency preserving witnesses of the consistencies between a total ideal and neighborhoods in a finite set.

**Lemma 5.8.** *At a type  $\rho$ , the statement (W) is equivalent to the following: Let  $\Gamma \in \text{Fin}_\rho$  and  $x \in G_\rho$ ; for all  $U \in \text{Con}_\Gamma$  with  $U \asymp_\rho x$  there exists a neighborhood  $N_{U, \Gamma, x} \in \text{Con}_\rho$  such that*

$$U \vdash_\rho N_{U, \Gamma, x} \subseteq x \wedge \bigvee_{U' \in \text{Con}_\Gamma} (U' \not\asymp_\rho U \rightarrow U' \not\asymp_\rho N_{U, \Gamma, x}). \quad (\text{W}')$$

*Proof.* Let  $\Gamma$  be a finite set,  $U$  some neighborhood of  $\Gamma$  and  $x$  a total ideal. Assume that (W) holds, and furthermore that  $U_1, \dots, U_m \in \text{Con}_\Gamma$  are all neighborhoods in  $\Gamma$  such that  $U_i \not\asymp_\rho U$  for  $i = 1, \dots, m$ . Then for each such  $i$  there is a neighborhood  $U_{0i} \in \text{Con}_\rho$

such that  $U_i \not\approx_\rho U_{0i}$  and  $U \vdash_\rho U_{0i} \subseteq x$ . Setting  $N_{U,\Gamma,x} := \bigcup_{i=1}^m U_{0i}$  we are done. In the other way around, let  $U$  and  $U'$  be two neighborhoods and  $x$  a total ideal, such that  $U' \not\approx_\rho U \approx_\rho x$ , and assume that (W') holds for all finite sets  $\Gamma$ , neighborhoods  $U \subseteq \Gamma$  and total ideals  $x$ . Setting  $U_0 := N_{U,U \cup U',x}$  we are done.  $\square$

So if  $\Gamma$  accepts a total ideal  $x$  at all, even if with a nontransitive neighborhood  $U$ , then it could be safely side extended to include a common part  $N_{U,\Gamma,x}$  of  $\overline{U}$  and  $x$ ; enriched in this way  $\Gamma$  would now accept  $x$  in the strong sense of inclusion. This is exactly what we need to exploit by taking the consistency closure of a higher-type total neighborhood, provided its list of arguments is omniceptive. But let us get to the details without further ado.

**Proposition 5.9** (Closure). *Let  $\rho$  and  $\sigma$  be types. If (O) and (W) hold in  $\rho$  and (C) holds in  $\sigma$  then (C) holds in  $\rho \rightarrow \sigma$ .*

*Proof.* Let  $T \in \text{Con}_{\rho \rightarrow \sigma}^g$  and  $x \in G_\rho$ . We show that  $\tilde{T}(x) \in G_\sigma$ . Based on Lemma 3.9, we may assume that  $L(T) \in \text{Fin}_\rho^o$  without harming generality. By (O) at  $\rho$  there exists a  $U^x \in \text{Con}_{L(T)}^{\text{cr}}$  such that  $U^x \approx_\rho x$ . By Lemma 4.8.4 we have  $T \cdot U^x \in \text{Con}_\sigma^g$ , and by (C) at  $\sigma$  we have  $\widetilde{T \cdot U^x} \in G_\sigma$ . So in order to show that  $\tilde{T}(x) \in G_\sigma$ , it suffices to show that  $\widetilde{T \cdot U^x} \subseteq \tilde{T}(x)$  and invoke Lemma 5.2.

Let then  $b \in \text{Tok}_\sigma$  be such that  $b \in \widetilde{T \cdot U^x}$ . This means that  $b \approx_\sigma T \cdot U^x$ . By Lemma 3.7.2 we have  $\langle U^x, b \rangle \approx_{\rho \rightarrow \sigma} T$ . By (W) at  $\rho$  and Lemma 5.8, there exists a neighborhood  $U_0^x := N_{U^x, L(T), x} \in \text{Con}_\rho$  such that

$$U^x \vdash_\rho U_0^x \subseteq x \wedge \bigvee_{U' \in \text{Con}_{L(T)}} (U' \not\approx_\rho U^x \rightarrow U' \not\approx_\rho U_0^x);$$

we have  $\langle U_0^x, b \rangle \approx_{\rho \rightarrow \sigma} T$ , because for every  $\langle U', b' \rangle \in T$  with  $U' \approx_\rho U_0^x$  it has to be  $U' \approx_\rho U^x$  from the above, therefore  $b \approx_\sigma b'$  follows by  $\langle U^x, b \rangle \approx_{\rho \rightarrow \sigma} T$ . We have found a  $U_b := U_0^x \in \text{Con}_\rho$  such that  $\langle U_b, b \rangle \approx_{\rho \rightarrow \sigma} T$  and  $U_b \subseteq x$ ; but this means by definition that  $b \in \tilde{T}(x)$ , and we are done.  $\square$

**Theorem 5.10** (Density). *Every type satisfies (O), (W), and (C), and in particular, every type is dense.*

*Proof.* It follows by a mutual induction over types by Lemma 5.5 and Propositions 5.4, 5.6, 5.7, and 5.9.  $\square$

As a closing remark, we should note that the witness which we provide is actually the *maximal* total extension of a given neighborhood, in the sense that if, for a type  $\rho$ ,  $U \in \text{Con}_\rho$  is some neighborhood,  $U^g \in \text{Con}_\rho^g$  is the witness provided by Theorem 4.7, and  $x \in G_\rho$  is such that  $U \subseteq x$ , then  $x \subseteq \widetilde{U^g}$ .

## 5.2 Nontotality of transitive ideals

In the same way as we did with finite totality and transitivity in Theorem 4.9, we would like to know if we can connect totality and transitivity on the level of ideals, and possibly obtain an explicit characterization of totality in terms of consistency. We show that this is not possible.

In [5, 6], an element  $x$  is defined to be *almost maximal* when  $y_1 \supseteq x \subseteq y_2$  implies  $y_1 \approx y_2$  for all  $y_1$  and  $y_2$ . At the same time, we call  $x$  *transitive* if  $y_1 \approx x \approx y_2$  implies

$y_1 \asymp y_2$  for all  $y_1, y_2$ . We also extend the bar notation for the deductive closure to ideals, in particular we write  $\overline{x \cup y}$  to mean the set  $\{a \mid \exists U \subseteq x \cup y \ U \vdash a\}$ . We immediately see the following.

**Lemma 5.11.** *At every type, an ideal is almost maximal if and only if it is transitive.*

*Proof.* That transitivity implies almost maximality is clear. To see the converse let  $x$  be almost maximal and  $y_1 \asymp x \asymp y_2$ . Then  $x \subseteq \overline{y_i \cup x}$  for each  $i$  and we get  $\overline{y_1 \cup x} \asymp \overline{y_2 \cup x}$  by almost maximality, which yields  $y_1 \asymp y_2$ .  $\square$

It is well known that in hierarchies over flat base types there exist functionals which are maximal—and therefore, by the above lemma, transitive—but not total, typical examples being various second order minimization operators [46, Example 8.3.2, Exercise 8.5.14]. In the nonflat setting we have counterexamples already at inductive base types, as we will see immediately. For the following we express the transitivity of  $x$  through tokens, similarly to Lemma 3.4.

**Proposition 5.12** (Total-transitivity). *At any type, total ideals are transitive. Conversely, there exist types with transitive ideals that are not total.*

*Proof.* At a base type  $\iota$ , let  $a_1, a_2$  be tokens and  $x$  a total ideal, such that  $a_1 \asymp_\iota x \asymp_\iota a_2$ . There exists a total neighborhood  $P$  with  $P \subseteq x$ , so the assumption yields  $a_1 \asymp_\iota P \asymp_\iota a_2$ , which implies  $a_1 \asymp_\iota a_2$  by the *finite* total transitivity of  $\iota$  (Proposition 4.3).

At a higher type  $\rho \rightarrow \sigma$ , let  $\langle U_1, b_1 \rangle, \langle U_2, b_2 \rangle$  be tokens and  $f$  be a total ideal, such that  $\langle U_1, b_1 \rangle \asymp_{\rho \rightarrow \sigma} f \asymp_{\rho \rightarrow \sigma} \langle U_2, b_2 \rangle$ . Assume furthermore that  $U_1 \asymp_\rho U_2$ . By Theorem 5.10 there exists a total ideal  $x : \rho$  such that  $U_1 \cup U_2 \subseteq x$ . Since  $f$  is itself total, the ideal  $f(x) : \sigma$  must also be total, and by the induction hypothesis at  $\sigma$  it must also be transitive. Now, applying all terms of the assumption to  $x$  we obtain  $b_1 \asymp_\sigma f(x) \asymp_\sigma b_2$ , which then yields  $b_1 \asymp_\sigma b_2$ .

For the converse, consider the ideal  $\infty = \{S^m * \mid m \geq 0\}$  of type  $\mathbb{N}$ .  $\square$

### 5.3 Noncontinuity of totalization

The witness for density that we have provided in section 5.1 is a mapping of the sort  $\text{tot} : \text{Con}_\rho \rightarrow \rho$ .<sup>6</sup> It is easy to see that this is not a “continuous” mapping—that is, it does not extend to an ideal of type  $\rho \rightarrow \rho$ —since it can not be expected to preserve consistency: consider the neighborhoods  $\{S*\}$  and  $\{SS*\}$  at type  $\mathbb{N}$ ; these are consistent with each other, but

$$\text{tot}(\{S*\}) \ni S0 \not\vdash_{\mathbb{N}} SS0 \in \text{tot}(\{SS*\}) .$$

This counterexample is general enough to convince us that this shortcoming is not particular to our witness.

**Lemma 5.13.** *There is no consistency-preserving mapping  $t : \text{Con}_{\mathbb{N}} \rightarrow \mathbb{N}$  such that  $U \subseteq t(U)$  and  $t(U) \in G_{\mathbb{N}}$  for all  $U \in \text{Con}_{\mathbb{N}}$ .*

*Proof.* If such a mapping existed it should be  $t(U_1) \asymp_{\mathbb{N}} t(U_2)$  for any two neighborhoods  $U_1, U_2 \subseteq \infty$ . Fixing such a  $U_1$  with  $t(U_1) = \{S^n 0\}$  for some  $n$  and setting  $U_2 := \{S^{n+1} * \}$  we get  $t(U_1) \not\vdash_{\mathbb{N}} t(U_2)$ , a contradiction.  $\square$

<sup>6</sup>Such mixed typings of terms appear often and naturally in considerations within information systems, and should be accounted for in a theory of partial computable functionals together with their approximations as in [20].

## 5.4 Separation

One of Berger’s key insights in [2], which permeates all subsequent approaches that our work is based upon (including our own), was that the notion of totality can be clarified if density is viewed together with an accompanying notion of “separation”: intuitively, a type  $\rho$  is considered to feature the *separation property*, if any two open sets of conflicting information can be told apart by a total “predicate” of type  $\rho \rightarrow \mathbb{B}$ . His argument proceeded by mutual induction for both properties of density and separation over all finite types. What we did instead in our mutual inductive arguments above was in effect to replace the notion of “separation of neighborhoods by infinite total ideals” by notions of “acceptance of total ideals by finite sets”. In our exposition separation follows as a simple corollary of density.

Following [44], and assuming the presence of the booleans in the type system, call a type  $\rho$  *separating* if

$$\forall_{U, U' \in \text{Con}_\rho} (U \not\#_\rho U' \rightarrow \exists_{f \in \tilde{G}_{\rho \rightarrow \mathbb{B}}} \langle U, \text{tt} \rangle \in f \ni \langle U', \text{ff} \rangle),$$

and *finitely separating* if

$$\forall_{U, U' \in \text{Con}_\rho} (U \not\#_\rho U' \rightarrow \exists_{T \in \text{Con}_{\rho \rightarrow \mathbb{B}}^g} \langle U, \text{tt} \rangle \succ_{\rho \rightarrow \mathbb{B}} T \succ_{\rho \rightarrow \mathbb{B}} \langle U', \text{ff} \rangle).$$

**Proposition 5.14** (Separation). *Every type is finitely separating, and consequently separating.*

*Proof.* If  $U$  and  $U'$  are inconsistent a  $\rho$ , then the finite set  $\{\langle U, \text{tt} \rangle, \langle U', \text{ff} \rangle\}$  is a neighborhood at  $\rho \rightarrow \mathbb{B}$ , and by Theorem 4.7 there will exist some  $T \in \text{Con}_{\rho \rightarrow \mathbb{B}}^g$  which side extends it. Consequently, by Theorem 5.10 the total ideal  $\tilde{T}$  will extend it.  $\square$

## 6 Notes

We gave a new, bottom-up proof of the Kreisel density theorem for finite types interpreted over nonflat inductive base types given as algebras by constructors. We introduced a notion of *totality for neighborhoods* and proved a *finite density theorem*, which states that, given a neighborhood, one may first totalize it in an explicitly finitary way to obtain a total neighborhood. Kreisel density is obtained by extending this total neighborhood to a total ideal by means of consistency; the resulting ideal, though generated by a compact element, is the maximal totalization of the given neighborhood. Here we gather notes on the above, on related literature, and on future work.

### The density theorem in the literature

As already pointed out in the introduction, the density problem was addressed for the first time by Kreisel [25] and also Kleene [24]. In his phd thesis [2], Ulrich Berger recast and solved the density problem within domain theory, generalizing results of Yuri Ershov [13, 14] and paralleling work of Dag Normann [30]—see [3, 46] for an account in English. A proof which does not thematize separation is given by Dag Normann in [33], while a modern approach from a viewpoint of an all-encompassing theory of higher-type computability can be found in the recent volume by John Longley and Dag Normann [26]. The density theorem is a fundamental result with several deep and

far-reaching applications, like the choice theorem [25, 3, 42], Kreisel’s representation theorem [25, 29, 32], a generalized Kreisel–Lacombe–Shoenfield theorem [3], Normann’s theorem [34, 35, 38], and Escardó’s theory of exhaustive search [15, 16], as well as extensions and generalizations, for example to dependent and universe domains [4], to Scott’s equilogical spaces [1], or even to an account of totality independently of density [31]—see also [5, 33, 6, 36]. It would be natural to seek among these studies for ones that would benefit from the possibility of explicitly finitary totalization. Existence of such cases would further justify the extension of the results presented here to richer type systems, starting with the one adopted in [44], and possibly moving on to the type systems covered in [4].

### Related work

The problem of finding a proof of density theorem “by compacts” occurred to the author back in the early 2011, and since then tackling it has primarily provided an incentive to develop the theory of nonflat information systems for semantics (see [23] for examples of collateral results). A partial result in the direction of finite witnesses for density was presented in [21], where, in contrast to the present approach, it was shown that one may first prove a version of finite separation at every type and then use this as a lemma to prove density (a version of our Proposition 5.7 also appears there); that approach provided a satisfactory finitary explanation of separation but not of totalization. Meanwhile, an alternative bottom-up approach to the density theorem, which grew independently but turned out to be similar in spirit to ours, was carried out by Davide Rinaldi in [39]. Rinaldi offers a nonflat semantics which is topological rather than domain-theoretic: he uses certain formal topologies [41], for which he proves that they are equivalent to *unary information systems*; these are information systems where in addition neighborhoods always have eigentokens, that is, for every  $U \in \text{Con}$  there exists some  $a \in \text{Tok}$  such that  $U \sim \{a\}$ . In our setting this is true of base types, but not of higher types. To adapt Rinaldi’s semantics in a way that clearly matches broader categories of information systems than just the unary ones, and look at a formal-topological proof of density by compacts anew, would not only be instructive, but it could also provide a more succinct and elegant proof.

### Towards a common study of totality and cototality

Recently, “cototal ideals”, that is, total ideals together with infinities like  $\infty$  at type  $\mathbb{N}$ , have been used to model stream-like objects at base types arising from initial algebras, offering an alternative to versions of semantics simultaneously based on initial algebras and final coalgebras [40, 18, 17]; for this line of work, rooted in [7, 8, 11], see [9, 44, 27, 28, 10]. In view of the mismatch between transitivity and totality in a nonflat setting which we described in section 5.2, it looks like a refinement is possible, where totality should feature an increased degree of finiteness and should be studied hand in hand with an appropriate notion of cototality: beside more or less obvious differences of the two at base types (based on the proof of Lemma 5.13, for example, one could expect continuous “cototalizations” to exist), their interplay at higher types remains terra incognita at the time of this writing.

## Acknowledgments

Thanks to Matthias Hofer for the feedback, Davide Rinaldi for the stimulating exchange of information, Parménides García Cornejo and Kenji Miyamoto for hearing me out, Apostolos Damialis for his advice on the notation, and to Dirk-André Deckert for all the backup. Special thanks to the referees for their substantial feedback: they shook me out of some misconceptions, posed insightful questions, and helped make the text a little more readable.

## References

- [1] Andrej Bauer and Lars Birkedal. Continuous functionals of dependent types and equilogical spaces. In *Computer science logic. 14th international workshop, CSL 2000. Annual conference of the EACSL, Fischbachau, Germany, August 21–26, 2000. Proceedings*, pages 202–216. Berlin: Springer, 2000.
- [2] Ulrich Berger. Totale Objekte und Mengen in der Bereichstheorie. (Total objects and sets in domain theory.). München: Univ. München, Fak. f. Mathematik. ii, 122 S. (1990)., 1990.
- [3] Ulrich Berger. Total sets and objects in domain theory. *Ann. Pure Appl. Logic*, 60(2):91–117, 1993.
- [4] Ulrich Berger. Continuous functionals of dependent and transfinite types. In *Models and computability. Invited papers from the Logic colloquium '97, European meeting of the Association for Symbolic Logic, Leeds, UK, July 6–13, 1997*, pages 1–22. Cambridge: Cambridge University Press, 1999.
- [5] Ulrich Berger. Effectivity and density in domains: A survey. In *A tutorial workshop on realizability semantics and applications. A workshop associated to the federated logic conference, Trento, Italy, June 30 – July 1, 1999*, page 13. Amsterdam: Elsevier, 1999.
- [6] Ulrich Berger. Computability and totality in domains. *Math. Struct. Comput. Sci.*, 12(3):281–294, 2002.
- [7] Ulrich Berger. Realisability and adequacy for (co)induction. In *6th international conference on computability and complexity in analysis (CCA'09). Proceedings of the international conference, August 18–22, 2009, Ljubljana, Slovenia*, page 12. Wadern: Schloss Dagstuhl – Leibniz Zentrum für Informatik, 2009.
- [8] Ulrich Berger. From coinductive proofs to exact real arithmetic: theory and applications. *Log. Methods Comput. Sci.*, 7(1):24, 2011.
- [9] Ulrich Berger, Kenji Miyamoto, Helmut Schwichtenberg, and Monika Seisenberger. Minlog - A tool for program extraction supporting algebras and coalgebras. In *Algebra and Coalgebra in Computer Science - 4th International Conference, CALCO 2011, Winchester, UK, August 30 - September 2, 2011. Proceedings*, pages 393–399, 2011.
- [10] Ulrich Berger, Kenji Miyamoto, Helmut Schwichtenberg, and Hideki Tsuiki. Logic for gray-code computation. In *Concepts of Proof in Mathematics, Philosophy, and Computer Science*. De Gruyter, 2016.
- [11] Ulrich Berger and Monika Seisenberger. Proofs, programs, processes. *Theory Comput. Syst.*, 51(3):313–329, 2012.
- [12] Yuri L. Ershov. Maximal and everywhere-defined functionals. *Algebra Logic*, 13:210–225, 1975.
- [13] Yuri L. Ershov. Theorie der Numerierungen. II. (Theory of numberings. II.) Übersetzung aus dem Russischen: H.-D. Hecker. Wissenschaftliche Redaktion: G. Asser. Berlin: VEB Deutscher Verlag der Wissenschaften. M 15.00 (1976). Sonderdruck aus *Z. math. Logik Grundle. Math.* 21, 473-584 (1975)., 1975.
- [14] Yuri L. Ershov. Model  $\mathbb{C}$  of partial continuous functionals. *Logic colloquium 76, Proc. Conf., Oxford 1976, Stud. Logic Found. Math., Vol. 87, 455-467 (1977).*, 1977.
- [15] Martín H. Escardó. Infinite sets that admit fast exhaustive search. In *Proceedings of the 22Nd Annual IEEE Symposium on Logic in Computer Science, LICS '07*, pages 443–452, Washington, DC, USA, 2007. IEEE Computer Society.
- [16] Martín H. Escardó. Exhaustible sets in higher-type computation. *Log. Methods Comput. Sci.*, 4(3):paper 3, 37, 2008.
- [17] Neil Ghani, Peter G. Hancock, and Dirk Pattinson. Continuous functions on final coalgebras. In *Proceedings of the 25th conference on the mathematical foundations of programming semantics (MFPS 2009), Oxford, UK, April 3–7, 2009*, pages 3–18. Amsterdam: Elsevier, 2009.

- [18] Peter G. Hancock, Neil Ghani, and Dirk Pattinson. Representations of stream processors using nested fixed points. *Log. Methods Comput. Sci.*, 5(3):17, 2009.
- [19] Simon Huber. On the computational content of choice axioms. Master’s thesis, Mathematisches Institut, LMU, 2010.
- [20] Simon Huber, Basil A. Karádaís, and Helmut Schwichtenberg. Towards a formal theory of computability. In *Ways of Proof Theory (Pohler’s Festschrift)*, pages 257–282. Ontos Verlag, Frankfurt, 2010.
- [21] Basil A. Karádaís. *Towards an Arithmetic with Approximations*. PhD thesis, Mathematisches Institut, LMU, 2013.
- [22] Basil A. Karádaís. Atomicity, coherence of information, and point-free structures. *Ann. Pure Appl. Logic*, 167(9):753–769, 2016.
- [23] Basil A. Karádaís. Normal forms, linearity, and prime algebraicity over nonflat domains. *Mathematical Logic Quarterly*, to appear, 2018.
- [24] Stephen C. Kleene. Countable functionals. In *Constructivity in mathematics: Proceedings of the colloquium held at Amsterdam, 1957 (edited by A. Heyting)*, Studies in Logic and the Foundations of Mathematics, pages 81–100, Amsterdam, 1959. North-Holland Publishing Co.
- [25] Georg Kreisel. Interpretation of analysis by means of constructive functionals of finite types. In *Constructivity in mathematics: Proceedings of the colloquium held at Amsterdam, 1957 (edited by A. Heyting)*, Studies in Logic and the Foundations of Mathematics, pages 101–128, Amsterdam, 1959. North-Holland Publishing Co.
- [26] John Longley and Dag Normann. *Higher-order computability*. Berlin: Springer, 2015.
- [27] Kenji Miyamoto, Fredrik Nordvall Forsberg, and Helmut Schwichtenberg. Program extraction from nested definitions. In *Interactive Theorem Proving - 4th International Conference, ITP 2013, Rennes, France, July 22-26, 2013. Proceedings*, pages 370–385, 2013.
- [28] Kenji Miyamoto and Helmut Schwichtenberg. Program extraction in exact real arithmetic. *Math. Struct. Comput. Sci.*, 25(8):1692–1704, 2015.
- [29] Dag Normann. Countable functionals and the projective hierarchy. *J. Symb. Log.*, 46:209–215, 1981.
- [30] Dag Normann. Kleene-spaces. Logic colloq. ’88, Proc. Colloq., Padova/Italy 1988, Stud. Logic Found. Math. 127, 91-109 (1989), 1989.
- [31] Dag Normann. A hierarchy of domains with totality, but without density. In *Computability, enumerability, unsolvability. Directions in recursion theory*, pages 233–257. Cambridge: Cambridge University Press, 1996.
- [32] Dag Normann. Closing the gap between the continuous functionals and recursion in <sup>3</sup>E. *Arch. Math. Logic*, 36(4-5):269–287, 1997.
- [33] Dag Normann. The continuous functionals. In *Handbook of computability theory*, pages 251–275. Amsterdam: Elsevier, 1999.
- [34] Dag Normann. Computability over the partial continuous functionals. *J. Symbolic Logic*, 65(3):1133–1142, 2000.
- [35] Dag Normann. The Cook-Berger problem. A guide to the solution. In *Domains IV. Workshop, Haus Humboldtstein, Remagen-Rolandseck, Germany, October 2–4, 1998*, page 9. Amsterdam: Elsevier, 2000.
- [36] Dag Normann. Applications of the Kleene-Kreisel density theorem to theoretical computer science. In *New computational paradigms. Changing conceptions of what is computable.*, pages 119–138. New York, NY: Springer, 2008.
- [37] Gordon D. Plotkin.  $T^0$  as a universal domain. *J. Comput. System Sci.*, 17(2):209–236, 1978.
- [38] Gordon D. Plotkin. Full abstraction, totality and PCF. *Math. Struct. Comput. Sci.*, 9(1):1–20, 1999.
- [39] Davide Rinaldi. *Formal methods in the theories of rings and domains*. PhD thesis, Mathematisches Institut, LMU, 2014.
- [40] Jan J.M.M. Rutten. Universal coalgebra: A theory of systems. *Theor. Comput. Sci.*, 249(1):3–80, 2000.
- [41] Giovanni Sambin. Intuitionistic formal spaces — a first communication. In *Mathematical logic and its applications (Druzhba, 1986)*, pages 187–204. Plenum, New York, 1987.
- [42] Helmut Schwichtenberg. Density and choice for total continuous functionals. In *Kreiseliana: about and around Georg Kreisel*, pages 335–362. Wellesley, MA: A K Peters, 1996.

- [43] Helmut Schwichtenberg. Recursion on the partial continuous functionals. In C. Dimitracopoulos, L. Newelski, D. Normann, and J. Steel, editors, *Logic Colloquium '05*, volume 28 of *Lecture Notes in Logic*, pages 173–201. Association for Symbolic Logic, 2007.
- [44] Helmut Schwichtenberg and Stanley S. Wainer. *Proofs and computations*. Perspectives in Logic. Cambridge University Press, Cambridge, 2012.
- [45] Dana S. Scott. Domains for denotational semantics. In *Automata, languages and programming (Aarhus, 1982)*, volume 140 of *Lecture Notes in Comput. Sci.*, pages 577–613. Springer, Berlin, 1982.
- [46] Viggo Stoltenberg-Hansen, Ingrid Lindström, and Edward R. Griffor. *Mathematical theory of domains*, volume 22 of *Cambridge Tracts in Theoretical Computer Science*. Cambridge University Press, 1994.
- [47] Glynn Winskel and Kim G. Larsen. Using information systems to solve recursive domain equations effectively. In *Semantics of Data Types*, pages 109–129, 1984.