

Mathematics of magic angles in twisted bilayer graphene


Herrsching, Mathematical results of many-body quantum systems


Simon Becker

June 10, 2022

NYU | COURANT








Based on joint work with

2020:, **Mark Embree, Jens Wittsten, Maciej Zworski**

2022: **Tristan Humbert, Maciej Zworski**



Weird behaviour of non-selfadjoint operators

Weird behaviour of non-selfadjoint operators

Seeley 85:

$$P(\alpha) = e^{ix} D_x + \alpha e^{ix}, \quad x \in \mathbb{R}/2\pi\mathbb{Z}, \quad D_x = \frac{1}{i} \partial_x$$

Weird behaviour of non-selfadjoint operators

Seeley 85:

$$P(\alpha) = e^{ix} D_x + \alpha e^{ix}, \quad x \in \mathbb{R}/2\pi\mathbb{Z}, \quad D_x = \frac{1}{i} \partial_x$$

$$\sigma(P(\alpha))$$

Weird behaviour of non-selfadjoint operators

Seeley 85:

$$P(\alpha) = e^{ix} D_x + \alpha e^{ix}, \quad x \in \mathbb{R}/2\pi\mathbb{Z}, \quad D_x = \frac{1}{i} \partial_x$$

$$\sigma(P(\alpha)) = \begin{cases} \mathbb{C}, & \alpha \in \mathbb{Z} \\ \emptyset, & \alpha \notin \mathbb{Z} \end{cases}$$

Weird behaviour of non-selfadjoint operators

Seeley 85:

$$P(\alpha) = e^{ix} D_x + \alpha e^{ix}, \quad x \in \mathbb{R}/2\pi\mathbb{Z}, \quad D_x = \frac{1}{i} \partial_x$$

$$\sigma(P(\alpha)) = \begin{cases} \mathbb{C}, & \alpha \in \mathbb{Z} \\ \emptyset, & \alpha \notin \mathbb{Z} \end{cases}$$

For all $\alpha \in \mathbb{C}$, $P(\alpha)$ is an **elliptic Fredholm operator of index 0** !

Weird behaviour of non-selfadjoint operators

Seeley 85:


$$P(\alpha) = e^{ix} D_x + \alpha e^{ix}, \quad x \in \mathbb{R}/2\pi\mathbb{Z}, \quad D_x = \frac{1}{i} \partial_x$$

$$\sigma(P(\alpha)) = \begin{cases} \mathbb{C}, & \alpha \in \mathbb{Z} \\ \emptyset, & \alpha \notin \mathbb{Z} \end{cases}$$


For all $\alpha \in \mathbb{C}$, $P(\alpha)$ is an **elliptic Fredholm operator of index 0** !

E. Brian Davies 2007: *Studying non-self-adjoint operators is like being a vet rather than a doctor: one has to require a much wider range of knowledge, and accept that one cannot expect to have as high a rate of success when confronted with particular cases.*


Motivation: bilayer graphene




Motivation: bilayer graphene




MacGyver in the physics lab



Motivation: bilayer graphene




MacGyver in the physics lab




Geim–Novoselov '04

Motivation: bilayer graphene



MacGyver in the physics lab





Geim–Novoselov '04




192 USD for 500 mg

Motivation: bilayer graphene





Geim–Novoselov '04

192 USD for 500 mg




Motivation: bilayer graphene




Geim–Novoselov '04

192 USD for 500 mg




Tight-binding model-Twisted bilayer graphene



We define the discrete Laplacian on the honeycomb lattice $H : \ell^2(\mathbb{Z}^2; \mathbb{C}^2) \rightarrow \ell^2(\mathbb{Z}^2; \mathbb{C}^2)$ by

$$H = \begin{pmatrix} 0 & 1 + \tau_1^* + \tau_2^* \\ 1 + \tau_1 + \tau_2 & 0 \end{pmatrix},$$

Tight-binding model-Twisted bilayer graphene




We define the discrete Laplacian on the honeycomb lattice $H : \ell^2(\mathbb{Z}^2; \mathbb{C}^2) \rightarrow \ell^2(\mathbb{Z}^2; \mathbb{C}^2)$ by

$$H = \begin{pmatrix} 0 & 1 + \tau_1^* + \tau_2^* \\ 1 + \tau_1 + \tau_2 & 0 \end{pmatrix},$$

where $(\tau_i \psi)_n = \psi_{n-e_i}$.

Tight-binding model-Twisted bilayer graphene




We define the discrete Laplacian on the honeycomb lattice $H : \ell^2(\mathbb{Z}^2; \mathbb{C}^2) \rightarrow \ell^2(\mathbb{Z}^2; \mathbb{C}^2)$ by

$$H = \begin{pmatrix} 0 & 1 + \tau_1^* + \tau_2^* \\ 1 + \tau_1 + \tau_2 & 0 \end{pmatrix},$$

where $(\tau_i \psi)_n = \psi_{n-e_i}$. The Hamiltonian describing twisted bilayer graphene is then given by

$$\mathcal{H}_\theta = \begin{pmatrix} H & W_\theta^* \\ W_\theta & H \end{pmatrix} : \ell^2(\mathbb{Z}^2; \mathbb{C}^4) \rightarrow \ell^2(\mathbb{Z}^2; \mathbb{C}^4),$$

Tight-binding model-Twisted bilayer graphene



We define the discrete Laplacian on the honeycomb lattice $H : \ell^2(\mathbb{Z}^2; \mathbb{C}^2) \rightarrow \ell^2(\mathbb{Z}^2; \mathbb{C}^2)$ by

$$H = \begin{pmatrix} 0 & 1 + \tau_1^* + \tau_2^* \\ 1 + \tau_1 + \tau_2 & 0 \end{pmatrix},$$

where $(\tau_i \psi)_n = \psi_{n-e_i}$. The Hamiltonian describing twisted bilayer graphene is then given by

$$\mathcal{H}_\theta = \begin{pmatrix} H & W_\theta^* \\ W_\theta & H \end{pmatrix} : \ell^2(\mathbb{Z}^2; \mathbb{C}^4) \rightarrow \ell^2(\mathbb{Z}^2; \mathbb{C}^4),$$

where

$$(W_\theta \psi)_n = (w * \psi)(R_\theta n) := \sum_{m \in \mathbb{Z}^2} w(R_\theta n - m) \psi_m \text{ with } w \in \mathcal{S}(\mathbb{R}^2; \mathbb{C}^{2 \times 2}).$$

The Bistritzer-MacDonald Hamiltonian

Massatt-Carr'20 showed that close to zero energy the model is effectively described by:

The Hamiltonian of two non-interacting sheets of graphene is

$$H = \begin{pmatrix} H_D & 0 \\ 0 & H_D \end{pmatrix} \text{ with } H_D = \begin{pmatrix} 0 & 2D_{\bar{z}} \\ 2D_z & 0 \end{pmatrix}.$$

The Bistritzer-MacDonald Hamiltonian

Massatt-Carr'20 showed that close to zero energy the model is effectively described by:

The Hamiltonian of two non-interacting sheets of graphene is

$$H = \begin{pmatrix} H_D & 0 \\ 0 & H_D \end{pmatrix} \text{ with } H_D = \begin{pmatrix} 0 & 2D_{\bar{z}} \\ 2D_z & 0 \end{pmatrix}.$$

The interacting between sheets with twisting angle θ is through tunnelling with Γ -periodic potentials

$$V_{\text{tun}} = \begin{pmatrix} V(\theta z) & \overline{U(\theta z)} \\ U(-\theta z) & V(\theta) \end{pmatrix}.$$

The Bistritzer-MacDonald Hamiltonian

Massatt-Carr'20 showed that close to zero energy the model is effectively described by:

The Hamiltonian of two non-interacting sheets of graphene is

$$H = \begin{pmatrix} H_D & 0 \\ 0 & H_D \end{pmatrix} \text{ with } H_D = \begin{pmatrix} 0 & 2D_{\bar{z}} \\ 2D_z & 0 \end{pmatrix}.$$

The interacting between sheets with twisting angle θ is through tunnelling with Γ -periodic potentials

$$V_{\text{tun}} = \begin{pmatrix} V(\theta z) & \overline{U(\theta z)} \\ U(-\theta z) & V(\theta) \end{pmatrix}.$$

The Bistritzer-MacDonald Hamiltonian is

$$H_{\text{BM}} = \begin{pmatrix} H_D & V_{\text{tun}} \\ V_{\text{tun}}^* & H_D \end{pmatrix}.$$

The chiral model of TBG

The chiral model of TBG

PHYSICAL REVIEW LETTERS **122**, 106405 (2019)

Editors' Suggestion

Origin of Magic Angles in Twisted Bilayer Graphene

Grigory Tarnopolsky, Alex Jura Kruchkov, and Ashvin Vishwanath
Department of Physics, Harvard University, Cambridge, Massachusetts 02138, USA

Origin of Magic Angles in Twisted Bilayer Graphene

Grigory Tamopolsky, Alex Jura Kruchkov, and Ashvin Vishwanath
Department of Physics, Harvard University, Cambridge, Massachusetts 02138, USA

$$H(\alpha) := \begin{pmatrix} 0 & D(\alpha)^* \\ D(\alpha) & 0 \end{pmatrix}, \quad D(\alpha) := \begin{pmatrix} 2D_{\bar{z}} & \alpha U(z) \\ \alpha U(-z) & 2D_{\bar{z}} \end{pmatrix},$$

$$z = x_1 + ix_2, \quad D_{\bar{z}} := \frac{1}{2i}(\partial_{x_1} + i\partial_{x_2})$$

$$U(z) := \sum_{k=0}^2 \omega^k e^{\frac{1}{2}(z\bar{\omega}^k - \bar{z}\omega^k)}, \quad \omega := e^{2\pi i/3}.$$

Origin of Magic Angles in Twisted Bilayer Graphene

Grigory Tamopolsky, Alex Jura Kruchkov, and Ashvin Vishwanath
 Department of Physics, Harvard University, Cambridge, Massachusetts 02138, USA

$$H(\alpha) := \begin{pmatrix} 0 & D(\alpha)^* \\ D(\alpha) & 0 \end{pmatrix}, \quad D(\alpha) := \begin{pmatrix} 2D_{\bar{z}} & \alpha U(z) \\ \alpha U(-z) & 2D_{\bar{z}} \end{pmatrix},$$

$$z = x_1 + ix_2, \quad D_{\bar{z}} := \frac{1}{2i}(\partial_{x_1} + i\partial_{x_2})$$

$$U(z) := \sum_{k=0}^2 \omega^k e^{\frac{1}{2}(z\bar{\omega}^k - \bar{z}\omega^k)}, \quad \omega := e^{2\pi i/3}.$$

$$U(z + \frac{4}{3}\pi i \omega^\ell) = \bar{\omega} U(z), \quad U(\omega z) = \omega U(z), \quad \ell = 1, 2.$$

Origin of Magic Angles in Twisted Bilayer Graphene

Grigory Tamopolsky, Alex Jura Kruchkov, and Ashvin Vishwanath
 Department of Physics, Harvard University, Cambridge, Massachusetts 02138, USA

$$H(\alpha) := \begin{pmatrix} 0 & D(\alpha)^* \\ D(\alpha) & 0 \end{pmatrix}, \quad D(\alpha) := \begin{pmatrix} 2D_{\bar{z}} & \alpha U(z) \\ \alpha U(-z) & 2D_{\bar{z}} \end{pmatrix},$$

$$z = x_1 + ix_2, \quad D_{\bar{z}} := \frac{1}{2i}(\partial_{x_1} + i\partial_{x_2})$$

$$U(z) := \sum_{k=0}^2 \omega^k e^{\frac{1}{2}(z\bar{\omega}^k - \bar{z}\omega^k)}, \quad \omega := e^{2\pi i/3}.$$

$$U(z + \frac{4}{3}\pi i \omega^\ell) = \bar{\omega} U(z), \quad U(\omega z) = \omega U(z), \quad \ell = 1, 2.$$

This limit is very important also for many-body effects *cf.*
 Bernevig–Vishwanath–Zaletel et al.

Review of symmetries

$$D(\alpha) = \begin{pmatrix} 2D_{\bar{z}} & \alpha U(z) \\ \alpha U(-z) & 2D_{\bar{z}} \end{pmatrix}, \quad H(\alpha) = \begin{pmatrix} 0 & D^* \\ D & 0 \end{pmatrix}$$

Review of symmetries

$$D(\alpha) = \begin{pmatrix} 2D_{\bar{z}} & \alpha U(z) \\ \alpha U(-z) & 2D_{\bar{z}} \end{pmatrix}, \quad H(\alpha) = \begin{pmatrix} 0 & D^* \\ D & 0 \end{pmatrix}$$

$$\mathcal{L}_{\mathbf{a}} \mathbf{u} = \text{diag}(\omega^{a_1+a_2}, 1, \omega^{a_1+a_2}, 1) \mathbf{u}(z + \frac{4}{3}i\pi(\omega a_1 + \omega^2 a_2)), \quad \mathbf{a} \in \mathbb{Z}_3^2,$$

Review of symmetries

$$D(\alpha) = \begin{pmatrix} 2D_{\bar{z}} & \alpha U(z) \\ \alpha U(-z) & 2D_{\bar{z}} \end{pmatrix}, \quad H(\alpha) = \begin{pmatrix} 0 & D^* \\ D & 0 \end{pmatrix}$$

$$\mathcal{L}_{\mathbf{a}} \mathbf{u} = \text{diag}(\omega^{a_1+a_2}, 1, \omega^{a_1+a_2}, 1) \mathbf{u}(z + \frac{4}{3}i\pi(\omega a_1 + \omega^2 a_2)), \quad \mathbf{a} \in \mathbb{Z}_3^2,$$

$$\mathcal{C}^k \mathbf{u}(z) = \text{diag}(1, 1, \bar{\omega}^k, \bar{\omega}^k) \mathbf{u}(\omega^k z), \quad k \in \mathbb{Z}_3$$

Review of symmetries

$$D(\alpha) = \begin{pmatrix} 2D_{\bar{z}} & \alpha U(z) \\ \alpha U(-z) & 2D_{\bar{z}} \end{pmatrix}, \quad H(\alpha) = \begin{pmatrix} 0 & D^* \\ D & 0 \end{pmatrix}$$

$$\mathcal{L}_{\mathbf{a}}\mathbf{u} = \text{diag}(\omega^{a_1+a_2}, 1, \omega^{a_1+a_2}, 1)\mathbf{u}(z + \frac{4}{3}i\pi(\omega a_1 + \omega^2 a_2)), \quad \mathbf{a} \in \mathbb{Z}_3^2,$$

$$\mathcal{C}^k\mathbf{u}(z) = \text{diag}(1, 1, \bar{\omega}^k, \bar{\omega}^k)\mathbf{u}(\omega^k z), \quad k \in \mathbb{Z}_3$$

$$\mathcal{L}_{\mathbf{a}}H = H\mathcal{L}_{\mathbf{a}},$$

Review of symmetries

$$D(\alpha) = \begin{pmatrix} 2D_{\bar{z}} & \alpha U(z) \\ \alpha U(-z) & 2D_{\bar{z}} \end{pmatrix}, \quad H(\alpha) = \begin{pmatrix} 0 & D^* \\ D & 0 \end{pmatrix}$$

$$\mathcal{L}_{\mathbf{a}}\mathbf{u} = \text{diag}(\omega^{a_1+a_2}, 1, \omega^{a_1+a_2}, 1)\mathbf{u}(z + \frac{4}{3}i\pi(\omega a_1 + \omega^2 a_2)), \quad \mathbf{a} \in \mathbb{Z}_3^2,$$

$$\mathcal{C}^k\mathbf{u}(z) = \text{diag}(1, 1, \bar{\omega}^k, \bar{\omega}^k)\mathbf{u}(\omega^k z), \quad k \in \mathbb{Z}_3$$

$$\mathcal{L}_{\mathbf{a}}H = H\mathcal{L}_{\mathbf{a}}, \quad \mathcal{C}H = H\mathcal{C},$$

Review of symmetries

$$D(\alpha) = \begin{pmatrix} 2D_{\bar{z}} & \alpha U(z) \\ \alpha U(-z) & 2D_{\bar{z}} \end{pmatrix}, \quad H(\alpha) = \begin{pmatrix} 0 & D^* \\ D & 0 \end{pmatrix}$$

$$\mathcal{L}_{\mathbf{a}}\mathbf{u} = \text{diag}(\omega^{a_1+a_2}, 1, \omega^{a_1+a_2}, 1)\mathbf{u}(z + \frac{4}{3}i\pi(\omega a_1 + \omega^2 a_2)), \quad \mathbf{a} \in \mathbb{Z}_3^2,$$

$$\mathcal{C}^k\mathbf{u}(z) = \text{diag}(1, 1, \bar{\omega}^k, \bar{\omega}^k)\mathbf{u}(\omega^k z), \quad k \in \mathbb{Z}_3$$

$$\mathcal{L}_{\mathbf{a}}H = H\mathcal{L}_{\mathbf{a}}, \quad \mathcal{C}H = H\mathcal{C}, \quad \mathcal{C}\mathcal{L}_{\mathbf{a}} = \mathcal{L}_{M\mathbf{a}}\mathcal{C},$$

Review of symmetries

$$D(\alpha) = \begin{pmatrix} 2D_{\bar{z}} & \alpha U(z) \\ \alpha U(-z) & 2D_{\bar{z}} \end{pmatrix}, \quad H(\alpha) = \begin{pmatrix} 0 & D^* \\ D & 0 \end{pmatrix}$$

$$\mathcal{L}_{\mathbf{a}}\mathbf{u} = \text{diag}(\omega^{a_1+a_2}, 1, \omega^{a_1+a_2}, 1)\mathbf{u}(z + \frac{4}{3}i\pi(\omega a_1 + \omega^2 a_2)), \quad \mathbf{a} \in \mathbb{Z}_3^2,$$

$$\mathcal{C}^k\mathbf{u}(z) = \text{diag}(1, 1, \bar{\omega}^k, \bar{\omega}^k)\mathbf{u}(\omega^k z), \quad k \in \mathbb{Z}_3$$

$$\mathcal{L}_{\mathbf{a}}H = H\mathcal{L}_{\mathbf{a}}, \quad \mathcal{C}H = H\mathcal{C}, \quad \mathcal{C}\mathcal{L}_{\mathbf{a}} = \mathcal{L}_{M\mathbf{a}}\mathcal{C}, \quad M = \begin{pmatrix} 0 & -1 \\ 1 & -1 \end{pmatrix}.$$

Review of symmetries

$$D(\alpha) = \begin{pmatrix} 2D_{\bar{z}} & \alpha U(z) \\ \alpha U(-z) & 2D_{\bar{z}} \end{pmatrix}, \quad H(\alpha) = \begin{pmatrix} 0 & D^* \\ D & 0 \end{pmatrix}$$

$$\mathcal{L}_{\mathbf{a}} \mathbf{u} = \text{diag}(\omega^{a_1+a_2}, 1, \omega^{a_1+a_2}, 1) \mathbf{u}(z + \frac{4}{3}i\pi(\omega a_1 + \omega^2 a_2)), \quad \mathbf{a} \in \mathbb{Z}_3^2,$$

$$\mathcal{C}^k \mathbf{u}(z) = \text{diag}(1, 1, \bar{\omega}^k, \bar{\omega}^k) \mathbf{u}(\omega^k z), \quad k \in \mathbb{Z}_3$$

$$\mathcal{L}_{\mathbf{a}} H = H \mathcal{L}_{\mathbf{a}}, \quad \mathcal{C} H = H \mathcal{C}, \quad \mathcal{C} \mathcal{L}_{\mathbf{a}} = \mathcal{L}_{M\mathbf{a}} \mathcal{C}, \quad M = \begin{pmatrix} 0 & -1 \\ 1 & -1 \end{pmatrix}.$$

Decompose into irreducible representations of this **Heisenberg** group:

$$L^2(\mathbb{C}/\Gamma) = \bigoplus_{k,p \in \mathbb{Z}_3} L^2_{\rho_{k,p}}(\mathbb{C}/\Gamma; \mathbb{C}^2) \oplus L^2_{\rho_{(1,0)}}(\mathbb{C}/\Gamma; \mathbb{C}^2) \oplus L^2_{\rho_{(2,0)}}(\mathbb{C}/\Gamma; \mathbb{C}^2)$$

Review of symmetries

$$D(\alpha) = \begin{pmatrix} 2D_{\bar{z}} & \alpha U(z) \\ \alpha U(-z) & 2D_{\bar{z}} \end{pmatrix}, \quad H(\alpha) = \begin{pmatrix} 0 & D^* \\ D & 0 \end{pmatrix}$$

$$\mathcal{L}_{\mathbf{a}} \mathbf{u} = \text{diag}(\omega^{a_1+a_2}, 1, \omega^{a_1+a_2}, 1) \mathbf{u}(z + \frac{4}{3}i\pi(\omega a_1 + \omega^2 a_2)), \quad \mathbf{a} \in \mathbb{Z}_3^2,$$

$$\mathcal{C}^k \mathbf{u}(z) = \text{diag}(1, 1, \bar{\omega}^k, \bar{\omega}^k) \mathbf{u}(\omega^k z), \quad k \in \mathbb{Z}_3$$

$$\mathcal{L}_{\mathbf{a}} H = H \mathcal{L}_{\mathbf{a}}, \quad \mathcal{C} H = H \mathcal{C}, \quad \mathcal{C} \mathcal{L}_{\mathbf{a}} = \mathcal{L}_{M\mathbf{a}} \mathcal{C}, \quad M = \begin{pmatrix} 0 & -1 \\ 1 & -1 \end{pmatrix}.$$

Decompose into irreducible representations of this **Heisenberg** group:

$$L^2(\mathbb{C}/\Gamma) = \bigoplus_{k,p \in \mathbb{Z}_3} L^2_{\rho_{k,p}}(\mathbb{C}/\Gamma; \mathbb{C}^2) \oplus L^2_{\rho_{(1,0)}}(\mathbb{C}/\Gamma; \mathbb{C}^2) \oplus L^2_{\rho_{(2,0)}}(\mathbb{C}/\Gamma; \mathbb{C}^2)$$

$$\rho_{k,p} \longleftrightarrow \mathcal{L}_{\mathbf{a}} \equiv \omega^{k(a_1+a_2)}, \quad \mathcal{C} \equiv \bar{\omega}^p$$

Symmetry protected states

Symmetry protected states

$$\ker_{L^2(\mathbb{C}/\Gamma)} H(0) = \mathbb{C}^4, \quad \Gamma = 4i\pi(\omega a_1 + \omega^2 a_2)$$

Symmetry protected states

$$\ker_{L^2(\mathbb{C}/\Gamma)} H(0) = \mathbb{C}^4, \quad \Gamma = 4i\pi(\omega a_1 + \omega^2 a_2)$$

$$\mathbf{e}_1 \in L^2_{\rho_{1,0}}, \quad \mathbf{e}_2 \in L^2_{\rho_{0,0}}, \quad \mathbf{e}_3 \in L^2_{\rho_{1,1}}, \quad \mathbf{e}_4 \in L^2_{\rho_{0,1}}.$$

Symmetry protected states

$$\ker_{L^2(\mathbb{C}/\Gamma)} H(0) = \mathbb{C}^4, \quad \Gamma = 4i\pi(\omega a_1 + \omega^2 a_2)$$

$$\mathbf{e}_1 \in L^2_{\rho_{1,0}}, \quad \mathbf{e}_2 \in L^2_{\rho_{0,0}}, \quad \mathbf{e}_3 \in L^2_{\rho_{1,1}}, \quad \mathbf{e}_4 \in L^2_{\rho_{0,1}}.$$

$$H(\alpha) = -\mathcal{W} H(\alpha) \mathcal{W}^*, \quad \mathcal{W} := \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad \mathcal{W} \mathcal{C} = \mathcal{C} \mathcal{W}, \quad \mathcal{L}_{\mathbf{a}} \mathcal{W} = \mathcal{W} \mathcal{L}_{\mathbf{a}}$$

Symmetry protected states

$$\ker_{L^2(\mathbb{C}/\Gamma)} H(0) = \mathbb{C}^4, \quad \Gamma = 4i\pi(\omega a_1 + \omega^2 a_2)$$

$$\mathbf{e}_1 \in L^2_{\rho_{1,0}}, \quad \mathbf{e}_2 \in L^2_{\rho_{0,0}}, \quad \mathbf{e}_3 \in L^2_{\rho_{1,1}}, \quad \mathbf{e}_4 \in L^2_{\rho_{0,1}}.$$

$$H(\alpha) = -\mathcal{W} H(\alpha) \mathcal{W}^*, \quad \mathcal{W} := \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad \mathcal{W} \mathcal{C} = \mathcal{C} \mathcal{W}, \quad \mathcal{L}_{\mathbf{a}} \mathcal{W} = \mathcal{W} \mathcal{L}_{\mathbf{a}}$$

This implies that the spectrum of $H(\alpha)|_{L^2_{\rho_{k,\ell}}(\mathbb{C}/\Gamma)}$ is **even**


Symmetry protected states

$$\ker_{L^2(\mathbb{C}/\Gamma)} H(0) = \mathbb{C}^4, \quad \Gamma = 4i\pi(\omega a_1 + \omega^2 a_2)$$

$$\mathbf{e}_1 \in L^2_{\rho_{1,0}}, \quad \mathbf{e}_2 \in L^2_{\rho_{0,0}}, \quad \mathbf{e}_3 \in L^2_{\rho_{1,1}}, \quad \mathbf{e}_4 \in L^2_{\rho_{0,1}}.$$

$$H(\alpha) = -\mathcal{W} H(\alpha) \mathcal{W}^*, \quad \mathcal{W} := \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad \mathcal{W} \mathcal{C} = \mathcal{C} \mathcal{W}, \quad \mathcal{L}_{\mathbf{a}} \mathcal{W} = \mathcal{W} \mathcal{L}_{\mathbf{a}}$$

This implies that the spectrum of $H(\alpha)|_{L^2_{\rho_{k,\ell}}(\mathbb{C}/\Gamma)}$ is **even**




Symmetry protected states

$$\ker_{L^2(\mathbb{C}/\Gamma)} H(0) = \mathbb{C}^4, \quad \Gamma = 4i\pi(\omega a_1 + \omega^2 a_2)$$

$$\mathbf{e}_1 \in L^2_{\rho_{1,0}}, \quad \mathbf{e}_2 \in L^2_{\rho_{0,0}}, \quad \mathbf{e}_3 \in L^2_{\rho_{1,1}}, \quad \mathbf{e}_4 \in L^2_{\rho_{0,1}}.$$

$$H(\alpha) = -\mathcal{W} H(\alpha) \mathcal{W}^*, \quad \mathcal{W} := \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad \mathcal{W} \mathcal{C} = \mathcal{C} \mathcal{W}, \quad \mathcal{L}_{\mathbf{a}} \mathcal{W} = \mathcal{W} \mathcal{L}_{\mathbf{a}}$$

This implies that the spectrum of $H(\alpha)|_{L^2_{\rho_{k,\ell}}(\mathbb{C}/\Gamma)}$ is **even**



$$\dim \ker_{L^2(\mathbb{C}/\Gamma)}(H(\alpha)) \geq 4, \quad \dim \ker_{L^2(\mathbb{C}/\Gamma)}(D(\alpha)) \geq 2$$

Spectral characterization of flat bands

Spectral characterization of flat bands

$$H_{\mathbf{k}}(\alpha) := \begin{pmatrix} 0 & D(\alpha)^* - \bar{\mathbf{k}} \\ D(\alpha) - \mathbf{k} & 0 \end{pmatrix} : H_0^1(\mathbb{C}/\Gamma) \rightarrow L_0^2(\mathbb{C}/\Gamma),$$

$$L_0^2(\mathbb{C}/\Gamma) := \{\mathbf{u} \in L^2(\mathbb{C}/\Gamma) : \mathcal{L}_{\mathbf{a}}\mathbf{u} = \mathbf{u}, \mathbf{a} \in \frac{1}{3}\Gamma/\Gamma\}.$$

Bands: $\{E_j(\alpha, \mathbf{k})\}_{j \in \mathbb{Z} \setminus \{0\}} = \text{Spec}_{L_0^2} H_{\mathbf{k}}(\alpha)$, $E_{\pm 1}(\alpha, \mathbf{0}) = E_{\pm 1}(\alpha, -\mathbf{i}) = 0$.

Spectral characterization of flat bands

$$H_{\mathbf{k}}(\alpha) := \begin{pmatrix} 0 & D(\alpha)^* - \bar{\mathbf{k}} \\ D(\alpha) - \mathbf{k} & 0 \end{pmatrix} : H_0^1(\mathbb{C}/\Gamma) \rightarrow L_0^2(\mathbb{C}/\Gamma),$$

$$L_0^2(\mathbb{C}/\Gamma) := \{\mathbf{u} \in L^2(\mathbb{C}/\Gamma) : \mathcal{L}_{\mathbf{a}}\mathbf{u} = \mathbf{u}, \mathbf{a} \in \frac{1}{3}\Gamma/\Gamma\}.$$

Bands: $\{E_j(\alpha, \mathbf{k})\}_{j \in \mathbb{Z} \setminus \{0\}} = \text{Spec}_{L_0^2} H_{\mathbf{k}}(\alpha)$, $E_{\pm 1}(\alpha, \mathbf{0}) = E_{\pm 1}(\alpha, -\mathbf{i}) = 0$.

Flat band at 0 $\iff \text{Spec}_{L_0^2(\mathbb{C}/\Gamma)}(D(\alpha)) = \mathbb{C}$

Spectral characterization of flat bands

$$H_{\mathbf{k}}(\alpha) := \begin{pmatrix} 0 & D(\alpha)^* - \bar{\mathbf{k}} \\ D(\alpha) - \mathbf{k} & 0 \end{pmatrix} : H_0^1(\mathbb{C}/\Gamma) \rightarrow L_0^2(\mathbb{C}/\Gamma),$$

$$L_0^2(\mathbb{C}/\Gamma) := \{\mathbf{u} \in L^2(\mathbb{C}/\Gamma) : \mathcal{L}_{\mathbf{a}}\mathbf{u} = \mathbf{u}, \mathbf{a} \in \frac{1}{3}\Gamma/\Gamma\}.$$

Bands: $\{E_j(\alpha, \mathbf{k})\}_{j \in \mathbb{Z} \setminus \{0\}} = \text{Spec}_{L_0^2} H_{\mathbf{k}}(\alpha)$, $E_{\pm 1}(\alpha, \mathbf{0}) = E_{\pm 1}(\alpha, -\mathbf{i}) = 0$.

Flat band at 0 $\iff \text{Spec}_{L_0^2(\mathbb{C}/\Gamma)}(D(\alpha)) = \mathbb{C}$

Spectral characterization of flat bands

$$H_{\mathbf{k}}(\alpha) := \begin{pmatrix} 0 & D(\alpha)^* - \bar{\mathbf{k}} \\ D(\alpha) - \mathbf{k} & 0 \end{pmatrix} : H_0^1(\mathbb{C}/\Gamma) \rightarrow L_0^2(\mathbb{C}/\Gamma),$$

$$L_0^2(\mathbb{C}/\Gamma) := \{\mathbf{u} \in L^2(\mathbb{C}/\Gamma) : \mathcal{L}_{\mathbf{a}}\mathbf{u} = \mathbf{u}, \mathbf{a} \in \frac{1}{3}\Gamma/\Gamma\}.$$

Bands: $\{E_j(\alpha, \mathbf{k})\}_{j \in \mathbb{Z} \setminus \{0\}} = \text{Spec}_{L_0^2} H_{\mathbf{k}}(\alpha)$, $E_{\pm 1}(\alpha, \mathbf{0}) = E_{\pm 1}(\alpha, -\mathbf{i}) = 0$.

Flat band at 0 $\iff \text{Spec}_{L_0^2(\mathbb{C}/\Gamma)}(D(\alpha)) = \mathbb{C}$

Theorem (BEWZ '20) *There exists a discrete set $\mathcal{A} \subset \mathbb{C}$ such that*

$$\text{Spec}_{L_0^2(\mathbb{C}/\Gamma)} D(\alpha) = \begin{cases} 3\Gamma^* & \alpha \notin \mathcal{A} \\ \mathbb{C} & \alpha \in \mathcal{A}, \end{cases}$$

Spectral characterization of flat bands

$$H_{\mathbf{k}}(\alpha) := \begin{pmatrix} 0 & D(\alpha)^* - \bar{\mathbf{k}} \\ D(\alpha) - \mathbf{k} & 0 \end{pmatrix} : H_0^1(\mathbb{C}/\Gamma) \rightarrow L_0^2(\mathbb{C}/\Gamma),$$


$$L_0^2(\mathbb{C}/\Gamma) := \{\mathbf{u} \in L^2(\mathbb{C}/\Gamma) : \mathcal{L}_{\mathbf{a}}\mathbf{u} = \mathbf{u}, \mathbf{a} \in \frac{1}{3}\Gamma/\Gamma\}.$$

Bands: $\{E_j(\alpha, \mathbf{k})\}_{j \in \mathbb{Z} \setminus \{0\}} = \text{Spec}_{L_0^2} H_{\mathbf{k}}(\alpha)$, $E_{\pm 1}(\alpha, \mathbf{0}) = E_{\pm 1}(\alpha, -\mathbf{i}) = 0$.

Flat band at 0 $\iff \text{Spec}_{L_0^2(\mathbb{C}/\Gamma)}(D(\alpha)) = \mathbb{C}$

Theorem (BEWZ '20) *There exists a discrete set $\mathcal{A} \subset \mathbb{C}$ such that*

$$\text{Spec}_{L_0^2(\mathbb{C}/\Gamma)} D(\alpha) = \begin{cases} 3\Gamma^* & \alpha \notin \mathcal{A} \\ \mathbb{C} & \alpha \in \mathcal{A}, \end{cases}$$



Proof of $\text{Spec}_{L^2(\mathbb{C}/\Gamma)} D(\alpha) = \begin{cases} \Gamma^* & \alpha \notin \mathcal{A} \\ \mathbb{C} & \alpha \in \mathcal{A}, \end{cases}$

Proof of $\text{Spec}_{L^2(\mathbb{C}/\Gamma)} D(\alpha) = \begin{cases} \Gamma^* & \alpha \notin \mathcal{A} \\ \mathbb{C} & \alpha \in \mathcal{A}, \end{cases}$

$$D(\alpha) = 2D_{\bar{z}} + \alpha \mathbf{V} \implies \text{Spec } D(\alpha) = \text{Spec } D(\alpha) + \Gamma^*, \quad \text{Spec } D(0) = \Gamma^*$$

Proof of $\text{Spec}_{L^2(\mathbb{C}/\Gamma)} D(\alpha) = \begin{cases} \Gamma^* & \alpha \notin \mathcal{A} \\ \mathbb{C} & \alpha \in \mathcal{A}, \end{cases}$

$$D(\alpha) = 2D_{\bar{z}} + \alpha \mathbf{V} \implies \text{Spec } D(\alpha) = \text{Spec } D(\alpha) + \Gamma^*, \quad \text{Spec } D(0) = \Gamma^*$$

$$\forall \alpha \quad 0 \in \text{Spec } D(\alpha)$$

Proof of $\text{Spec}_{L^2(\mathbb{C}/\Gamma)} D(\alpha) = \begin{cases} \Gamma^* & \alpha \notin \mathcal{A} \\ \mathbb{C} & \alpha \in \mathcal{A}, \end{cases}$

$$D(\alpha) = 2D_{\bar{z}} + \alpha \mathbf{V} \implies \text{Spec } D(\alpha) = \text{Spec } D(\alpha) + \Gamma^*, \quad \text{Spec } D(0) = \Gamma^*$$

$$\forall \alpha \neq 0 \in \text{Spec } D(\alpha) \implies \text{if } \text{Spec } D(\alpha) \text{ is discrete then } \text{Spec } D(\alpha) = \Gamma^*.$$

Proof of $\text{Spec}_{L^2(\mathbb{C}/\Gamma)} D(\alpha) = \begin{cases} \Gamma^* & \alpha \notin \mathcal{A} \\ \mathbb{C} & \alpha \in \mathcal{A}, \end{cases}$

$$D(\alpha) = 2D_{\bar{z}} + \alpha \mathbf{V} \implies \text{Spec } D(\alpha) = \text{Spec } D(\alpha) + \Gamma^*, \quad \text{Spec } D(0) = \Gamma^*$$

$$\forall \alpha \neq 0 \in \text{Spec } D(\alpha) \implies \text{if } \text{Spec } D(\alpha) \text{ is discrete then } \text{Spec } D(\alpha) = \Gamma^*.$$

$$\exists \mathbf{k} \quad (D(\alpha) - \mathbf{k})^{-1} : L^2 \rightarrow H^1 \iff \text{the spectrum is discrete}$$

Proof of $\text{Spec}_{L^2(\mathbb{C}/\Gamma)} D(\alpha) = \begin{cases} \Gamma^* & \alpha \notin \mathcal{A} \\ \mathbb{C} & \alpha \in \mathcal{A}, \end{cases}$

$$D(\alpha) = 2D_{\bar{z}} + \alpha \mathbf{V} \implies \text{Spec } D(\alpha) = \text{Spec } D(0) + \Gamma^*, \quad \text{Spec } D(0) = \Gamma^*$$

$$\forall \alpha \neq 0 \in \text{Spec } D(\alpha) \implies \text{if } \text{Spec } D(\alpha) \text{ is discrete then } \text{Spec } D(\alpha) = \Gamma^*.$$

$$\exists \mathbf{k} \quad (D(\alpha) - \mathbf{k})^{-1} : L^2 \rightarrow H^1 \iff \text{the spectrum is discrete}$$

$$(D(\alpha) - \mathbf{k})^{-1} = (I + \alpha T_{\mathbf{k}})^{-1} (D(0) - \mathbf{k})^{-1}, \quad T_{\mathbf{k}} := (D(0) - \mathbf{k})^{-1} \mathbf{V}, \quad \mathbf{k} \notin \Gamma^*$$

Proof of $\text{Spec}_{L^2(\mathbb{C}/\Gamma)} D(\alpha) = \begin{cases} \Gamma^* & \alpha \notin \mathcal{A} \\ \mathbb{C} & \alpha \in \mathcal{A}, \end{cases}$

$$D(\alpha) = 2D_{\bar{z}} + \alpha \mathbf{V} \implies \text{Spec } D(\alpha) = \text{Spec } D(0) + \Gamma^*, \quad \text{Spec } D(0) = \Gamma^*$$

$$\forall \alpha \neq 0 \in \text{Spec } D(\alpha) \implies \text{if } \text{Spec } D(\alpha) \text{ is discrete then } \text{Spec } D(\alpha) = \Gamma^*.$$

$$\exists \mathbf{k} \quad (D(\alpha) - \mathbf{k})^{-1} : L^2 \rightarrow H^1 \iff \text{the spectrum is discrete}$$

$$(D(\alpha) - \mathbf{k})^{-1} = (I + \alpha T_{\mathbf{k}})^{-1} (D(0) - \mathbf{k})^{-1}, \quad T_{\mathbf{k}} := (D(0) - \mathbf{k})^{-1} \mathbf{V}, \quad \mathbf{k} \notin \Gamma^*$$

$$\alpha \mapsto (I + \alpha T_{\mathbf{k}})^{-1} \quad \text{meromorphic with poles at } 1/\alpha \in \text{Spec}(T_{\mathbf{k}})$$

Proof of $\text{Spec}_{L^2(\mathbb{C}/\Gamma)} D(\alpha) = \begin{cases} \Gamma^* & \alpha \notin \mathcal{A} \\ \mathbb{C} & \alpha \in \mathcal{A}, \end{cases}$

$$D(\alpha) = 2D_{\bar{z}} + \alpha \mathbf{V} \implies \text{Spec } D(\alpha) = \text{Spec } D(0) + \Gamma^*, \quad \text{Spec } D(0) = \Gamma^*$$

$$\forall \alpha \neq 0 \in \text{Spec } D(\alpha) \implies \text{if } \text{Spec } D(\alpha) \text{ is discrete then } \text{Spec } D(\alpha) = \Gamma^*.$$

$$\exists \mathbf{k} \quad (D(\alpha) - \mathbf{k})^{-1} : L^2 \rightarrow H^1 \iff \text{the spectrum is discrete}$$

$$(D(\alpha) - \mathbf{k})^{-1} = (I + \alpha T_{\mathbf{k}})^{-1} (D(0) - \mathbf{k})^{-1}, \quad T_{\mathbf{k}} := (D(0) - \mathbf{k})^{-1} \mathbf{V}, \quad \mathbf{k} \notin \Gamma^*$$

$$\alpha \mapsto (I + \alpha T_{\mathbf{k}})^{-1} \text{ meromorphic with poles at } 1/\alpha \in \text{Spec}(T_{\mathbf{k}})$$

We obtained a spectral characterization of **magic angles** $\theta = 1/\alpha$:

$$\text{flat band} \iff \text{Spec}_{L^2(\mathbb{C}/\Gamma)} D(\alpha) = \mathbb{C} \iff 1/\alpha \in \text{Spec}(T_{\mathbf{k}})$$

Proof of $\text{Spec}_{L^2(\mathbb{C}/\Gamma)} D(\alpha) = \begin{cases} \Gamma^* & \alpha \notin \mathcal{A} \\ \mathbb{C} & \alpha \in \mathcal{A}, \end{cases}$

$$D(\alpha) = 2D_{\bar{z}} + \alpha \mathbf{V} \implies \text{Spec } D(\alpha) = \text{Spec } D(0) + \Gamma^*, \quad \text{Spec } D(0) = \Gamma^*$$

$$\forall \alpha \neq 0 \in \text{Spec } D(\alpha) \implies \text{if } \text{Spec } D(\alpha) \text{ is discrete then } \text{Spec } D(\alpha) = \Gamma^*.$$


$$\exists \mathbf{k} \quad (D(\alpha) - \mathbf{k})^{-1} : L^2 \rightarrow H^1 \iff \text{the spectrum is discrete}$$

$$(D(\alpha) - \mathbf{k})^{-1} = (I + \alpha T_{\mathbf{k}})^{-1} (D(0) - \mathbf{k})^{-1}, \quad T_{\mathbf{k}} := (D(0) - \mathbf{k})^{-1} \mathbf{V}, \quad \mathbf{k} \notin \Gamma^*$$

$$\alpha \mapsto (I + \alpha T_{\mathbf{k}})^{-1} \text{ meromorphic with poles at } 1/\alpha \in \text{Spec}(T_{\mathbf{k}})$$


We obtained a spectral characterization of **magic angles** $\theta = 1/\alpha$:

$$\text{flat band} \iff \text{Spec}_{L^2(\mathbb{C}/\Gamma)} D(\alpha) = \mathbb{C} \iff 1/\alpha \in \text{Spec}(T_{\mathbf{k}})$$




Flat bands

The bands are eigenvalues of $H_{\mathbf{k}}(\alpha)$ on $L_0^2(\mathbb{C}/\Gamma)$, $\mathbf{k} \in \mathbb{C}/3\Gamma^*$:



Flat bands

The bands are eigenvalues of $H_{\mathbf{k}}(\alpha)$ on $L_0^2(\mathbb{C}/\Gamma)$, $\mathbf{k} \in \mathbb{C}/3\Gamma^*$:



Theorem (BHZ '22; implicit in BEWZ '20)

$$\exists \mathbf{k} \notin 3\Gamma^* + \{\mathbf{0}, -\mathbf{i}\} \quad E_1(\alpha, \mathbf{k}) = 0 \implies \forall \mathbf{k} \quad E_1(\alpha, \mathbf{k}) = 0.$$


A curious structure of the first band

A curious structure of the first band

$$\mathbf{k} \mapsto \tilde{E}_1(\alpha, \mathbf{k}) / (\max_{\mathbf{k}} \tilde{E}_1(\alpha, \mathbf{k})), \quad 0.4 < \alpha < 0.6$$


A curious structure of the first band

$$\mathbf{k} \mapsto \tilde{E}_1(\alpha, \mathbf{k}) / (\max_{\mathbf{k}} \tilde{E}_1(\alpha, \mathbf{k})), \quad 0.4 < \alpha < 0.6$$



A curious structure of the first band


$$\mathbf{k} \mapsto \tilde{E}_1(\alpha, \mathbf{k}) / (\max_{\mathbf{k}} \tilde{E}_1(\alpha, \mathbf{k})), \quad 0.4 < \alpha < 0.6$$




Rescaled plots remain almost fixed at $\mathbf{k} \mapsto |U(-4\sqrt{3}\pi i\mathbf{k}/9)|$

Exponential squeezing of bands

Exponential squeezing of bands




Exponential squeezing of bands



Theorem. (BEWZ '20) *There exist $c_j > 0$ such that for all $\mathbf{k} \in \mathbb{C}$,*

$$|E_j(\alpha, \mathbf{k})| \leq c_0 e^{-c_1 \alpha}, \quad j \leq c_2 \alpha, \quad \alpha > 0.$$

Exponential squeezing of bands




Theorem. (BEWZ '20) *There exist $c_j > 0$ such that for all $\mathbf{k} \in \mathbb{C}$,*

$$|E_j(\alpha, \mathbf{k})| \leq c_0 e^{-c_1 \alpha}, \quad j \leq c_2 \alpha, \quad \alpha > 0.$$

In practice, $c_1 = 1$ and c_2 can be taken arbitrarily large

Exponential squeezing of bands




Theorem. (BEWZ '20) *There exist $c_j > 0$ such that for all $\mathbf{k} \in \mathbb{C}$,*

$$|E_j(\alpha, \mathbf{k})| \leq c_0 e^{-c_1 \alpha}, \quad j \leq c_2 \alpha, \quad \alpha > 0.$$

In practice, $c_1 = 1$ and c_2 can be taken arbitrarily large

Consequence of general results about **quasimodes** for semiclassical ($h = 1/\alpha$) non-normal operators:

Exponential squeezing of bands



Theorem. (BEWZ '20) *There exist $c_j > 0$ such that for all $\mathbf{k} \in \mathbb{C}$,*

$$|E_j(\alpha, \mathbf{k})| \leq c_0 e^{-c_1 \alpha}, \quad j \leq c_2 \alpha, \quad \alpha > 0.$$

In practice, $c_1 = 1$ and c_2 can be taken arbitrarily large

Consequence of general results about **quasimodes** for semiclassical ($h = 1/\alpha$) non-normal operators:

Hörmander'69, Sato-Kawai-Kashiwara'73...Dencker-Sjöstrand-Z'04

Exponential squeezing of bands via solvability of PDE

Exponential squeezing of bands via solvability of PDE

Lewy '57: $(\partial_{x_1} + i\partial_{x_2} - 2i(x_1 + ix_2)\partial_{x_3})u = f$ has **no** solution near any point $x \in \mathbb{R}^3$ for a **generic** $f \in C^\infty(\mathbb{R}^3)$

Exponential squeezing of bands via solvability of PDE

Lewy '57: $(\partial_{x_1} + i\partial_{x_2} - 2i(x_1 + ix_2)\partial_{x_3})u = f$ has **no** solution near any point $x \in \mathbb{R}^3$ for a **generic** $f \in C^\infty(\mathbb{R}^3)$

Hörmander '60: $P = \sum_{|\alpha| \leq m} a_\alpha(x) D_x^\alpha$, $p(x, \xi) := \sum_{|\alpha| = m} a_\alpha(x) \xi^\alpha$

Exponential squeezing of bands via solvability of PDE

Lewy '57: $(\partial_{x_1} + i\partial_{x_2} - 2i(x_1 + ix_2)\partial_{x_3})u = f$ has **no** solution near any point $x \in \mathbb{R}^3$ for a **generic** $f \in C^\infty(\mathbb{R}^3)$

Hörmander '60: $P = \sum_{|\alpha| \leq m} a_\alpha(x) D_x^\alpha$, $p(x, \xi) := \sum_{|\alpha|=m} a_\alpha(x) \xi^\alpha$

$$\exists (x_0, \xi_0), \xi_0 \neq 0, \quad p(x_0, \xi_0) = 0,$$

$$\{p, \bar{p}\} := \sum_{j=1}^n \partial_{\xi_j} p \partial_{x_j} \bar{p} - \partial_{\xi_j} \bar{p} \partial_{x_j} p|_{(x_0, \xi_0)} \neq 0$$

Exponential squeezing of bands via solvability of PDE

Lewy '57: $(\partial_{x_1} + i\partial_{x_2} - 2i(x_1 + ix_2)\partial_{x_3})u = f$ has **no** solution near any point $x \in \mathbb{R}^3$ for a **generic** $f \in C^\infty(\mathbb{R}^3)$

Hörmander '60: $P = \sum_{|\alpha| \leq m} a_\alpha(x) D_x^\alpha$, $p(x, \xi) := \sum_{|\alpha|=m} a_\alpha(x) \xi^\alpha$

$$\exists (x_0, \xi_0), \xi_0 \neq 0, \quad p(x_0, \xi_0) = 0,$$

$$\{p, \bar{p}\} := \sum_{j=1}^n \partial_{\xi_j} p \partial_{x_j} \bar{p} - \partial_{\xi_j} \bar{p} \partial_{x_j} p|_{(x_0, \xi_0)} \neq 0$$

$\implies Pu = f$ is **not** solvable near x_0 for a generic $f \in C^\infty$

Exponential squeezing of bands via solvability of PDE

Lewy '57: $(\partial_{x_1} + i\partial_{x_2} - 2i(x_1 + ix_2)\partial_{x_3})u = f$ has **no** solution near any point $x \in \mathbb{R}^3$ for a **generic** $f \in C^\infty(\mathbb{R}^3)$

Hörmander '60: $P = \sum_{|\alpha| \leq m} a_\alpha(x) D_x^\alpha$, $p(x, \xi) := \sum_{|\alpha|=m} a_\alpha(x) \xi^\alpha$

$$\exists (x_0, \xi_0), \xi_0 \neq 0, \quad p(x_0, \xi_0) = 0,$$

$$\{p, \bar{p}\} := \sum_{j=1}^n \partial_{\xi_j} p \partial_{x_j} \bar{p} - \partial_{\xi_j} \bar{p} \partial_{x_j} p|_{(x_0, \xi_0)} \neq 0$$

$\implies Pu = f$ is **not** solvable near x_0 for a generic $f \in C^\infty$

Essential step: $\exists u_h$ supported in $B(x_0, h^{\frac{1}{2}-})$ such that

$$\forall N \exists C_N \quad \|Pu_h\|_{L^2} \leq C_N h^N, \quad \|u_h\|_{L^2} = 1$$

Sato–Kawai–Kashiwara '73, Dencker–Sjöstrand–Z '04: if a_α 's are analytic functions then

$$\exists c > 0 \quad \|Pu_h\|_{L^2} \leq e^{-c/h}, \quad \|u_h\|_{L^2} = 1$$


$$\text{Spec}_{L^2(\mathbb{C}/\Gamma)} D(\alpha) = \begin{cases} \Gamma^* & \alpha \notin \mathcal{A} \\ \mathbb{C} & \alpha \in \mathcal{A}, \end{cases}$$

$$\text{Spec}_{L^2(\mathbb{C}/\Gamma)} D(\alpha) = \begin{cases} \Gamma^* & \alpha \notin \mathcal{A} \\ \mathbb{C} & \alpha \in \mathcal{A}, \end{cases}$$

flat band at $\alpha \iff \text{Spec}_{L^2(\mathbb{C}/\Gamma)} D(\alpha) = \mathbb{C} \iff 1/\alpha \in \text{Spec}(T_{\mathbf{k}})$

$$\text{Spec}_{L^2(\mathbb{C}/\Gamma)} D(\alpha) = \begin{cases} \Gamma^* & \alpha \notin \mathcal{A} \\ \mathbb{C} & \alpha \in \mathcal{A}, \end{cases}$$


flat band at $\alpha \iff \text{Spec}_{L^2(\mathbb{C}/\Gamma)} D(\alpha) = \mathbb{C} \iff 1/\alpha \in \text{Spec}(T_{\mathbf{k}})$



We did **not** prove that $\mathcal{A} \cap \mathbb{R}_+ \neq \emptyset$.

$$\text{Spec}_{L^2(\mathbb{C}/\Gamma)} D(\alpha) = \begin{cases} \Gamma^* & \alpha \notin \mathcal{A} \\ \mathbb{C} & \alpha \in \mathcal{A}, \end{cases}$$


flat band at $\alpha \iff \text{Spec}_{L^2(\mathbb{C}/\Gamma)} D(\alpha) = \mathbb{C} \iff 1/\alpha \in \text{Spec}(T_{\mathbf{k}})$



We did **not** prove that $\mathcal{A} \cap \mathbb{R}_+ \neq \emptyset$. However, $\mathcal{A} \neq \emptyset$ **BEWZ** '21:

$$\text{Spec}_{L^2(\mathbb{C}/\Gamma)} D(\alpha) = \begin{cases} \Gamma^* & \alpha \notin \mathcal{A} \\ \mathbb{C} & \alpha \in \mathcal{A}, \end{cases}$$

flat band at $\alpha \iff \text{Spec}_{L^2(\mathbb{C}/\Gamma)} D(\alpha) = \mathbb{C} \iff 1/\alpha \in \text{Spec}(T_{\mathbf{k}})$




We did **not** prove that $\mathcal{A} \cap \mathbb{R}_+ \neq \emptyset$. However, $\mathcal{A} \neq \emptyset$ **BEWZ** '21:

$$\sum_{\alpha \in \mathcal{A}} \alpha^{-4} = \text{tr } T_{\mathbf{k}}^4 = \frac{72\pi}{\sqrt{3}},$$

$$\text{Spec}_{L^2(\mathbb{C}/\Gamma)} D(\alpha) = \begin{cases} \Gamma^* & \alpha \notin \mathcal{A} \\ \mathbb{C} & \alpha \in \mathcal{A}, \end{cases}$$

flat band at $\alpha \iff \text{Spec}_{L^2(\mathbb{C}/\Gamma)} D(\alpha) = \mathbb{C} \iff 1/\alpha \in \text{Spec}(T_{\mathbf{k}})$




We did **not** prove that $\mathcal{A} \cap \mathbb{R}_+ \neq \emptyset$. However, $\mathcal{A} \neq \emptyset$ **BEWZ** '21:

$$\sum_{\alpha \in \mathcal{A}} \alpha^{-4} = \text{tr } T_{\mathbf{k}}^4 = \frac{72\pi}{\sqrt{3}}, \quad \text{combinatorics} + \wp \text{ function}$$

$$\text{Spec}_{L^2(\mathbb{C}/\Gamma)} D(\alpha) = \begin{cases} \Gamma^* & \alpha \notin \mathcal{A} \\ \mathbb{C} & \alpha \in \mathcal{A}, \end{cases}$$

flat band at $\alpha \iff \text{Spec}_{L^2(\mathbb{C}/\Gamma)} D(\alpha) = \mathbb{C} \iff 1/\alpha \in \text{Spec}(T_{\mathbf{k}})$




We did **not** prove that $\mathcal{A} \cap \mathbb{R}_+ \neq \emptyset$. However, $\mathcal{A} \neq \emptyset$ **BEWZ** '21:

$$\sum_{\alpha \in \mathcal{A}} \alpha^{-4} = \text{tr } T_{\mathbf{k}}^4 = \frac{72\pi}{\sqrt{3}}, \quad \text{combinatorics} + \wp \text{ function}$$

Luskin-Watson '21: computer assisted proof $|\mathcal{A} \cap \mathbb{R}_+| \geq 1$

$$\text{Spec}_{L^2(\mathbb{C}/\Gamma)} D(\alpha) = \begin{cases} \Gamma^* & \alpha \notin \mathcal{A} \\ \mathbb{C} & \alpha \in \mathcal{A}, \end{cases}$$

flat band at $\alpha \iff \text{Spec}_{L^2(\mathbb{C}/\Gamma)} D(\alpha) = \mathbb{C} \iff 1/\alpha \in \text{Spec}(T_{\mathbf{k}})$



We did **not** prove that $\mathcal{A} \cap \mathbb{R}_+ \neq \emptyset$. However, $\mathcal{A} \neq \emptyset$ **BEWZ** '21:

$$\sum_{\alpha \in \mathcal{A}} \alpha^{-4} = \text{tr } T_{\mathbf{k}}^4 = \frac{72\pi}{\sqrt{3}}, \quad \text{combinatorics} + \wp \text{ function}$$

Luskin–Watson '21: computer assisted proof $|\mathcal{A} \cap \mathbb{R}_+| \geq 1$

BEWZ '21 (motivated by **LW**): $|\mathcal{A} \cap \mathbb{R}_+| \geq 2$

$$\text{Spec}_{L^2(\mathbb{C}/\Gamma)} D(\alpha) = \begin{cases} \Gamma^* & \alpha \notin \mathcal{A} \\ \mathbb{C} & \alpha \in \mathcal{A}, \end{cases}$$

$$\text{Spec}_{L^2(\mathbb{C}/\Gamma)} D(\alpha) = \begin{cases} \Gamma^* & \alpha \notin \mathcal{A} \\ \mathbb{C} & \alpha \in \mathcal{A}, \end{cases}$$

Theorem (BHZ '22) For all $p > 1$

$$\sum_{\alpha \in \mathcal{A}} \alpha^{-2p} \in \frac{\pi}{\sqrt{3}} \mathbb{Q}$$

$$\text{Spec}_{L^2(\mathbb{C}/\Gamma)} D(\alpha) = \begin{cases} \Gamma^* & \alpha \notin \mathcal{A} \\ \mathbb{C} & \alpha \in \mathcal{A}, \end{cases}$$

Theorem (BHZ '22) For all $p > 1$

$$\sum_{\alpha \in \mathcal{A}} \alpha^{-2p} \in \frac{\pi}{\sqrt{3}} \mathbb{Q} \quad \text{and as a consequence } |\mathcal{A}| = \infty.$$

$$\text{Spec}_{L^2(\mathbb{C}/\Gamma)} D(\alpha) = \begin{cases} \Gamma^* & \alpha \notin \mathcal{A} \\ \mathbb{C} & \alpha \in \mathcal{A}, \end{cases}$$

Theorem (BHZ '22) For all $p > 1$

$$\sum_{\alpha \in \mathcal{A}} \alpha^{-2p} \in \frac{\pi}{\sqrt{3}} \mathbb{Q} \quad \text{and as a consequence } |\mathcal{A}| = \infty.$$

$$\sigma_p := \frac{1}{18} \text{tr } T_{\mathbf{k}}^{2p}, \quad F_{\mathbf{k}}(\alpha) := \det_2(I - \alpha^2 T_{\mathbf{k}}^2)$$

p	$\sqrt{3}\sigma_p/3^p\pi$
2	4/9
3	32/63
4	40/81

p	$\sqrt{3}\sigma_p/3^p\pi$
5	9560/20007
6	245120/527877
7	1957475168/4337177481

$$\text{Spec}_{L^2(\mathbb{C}/\Gamma)} D(\alpha) = \begin{cases} \Gamma^* & \alpha \notin \mathcal{A} \\ \mathbb{C} & \alpha \in \mathcal{A}, \end{cases}$$

Theorem (BHZ '22) For all $p > 1$

$$\sum_{\alpha \in \mathcal{A}} \alpha^{-2p} \in \frac{\pi}{\sqrt{3}} \mathbb{Q} \quad \text{and as a consequence } |\mathcal{A}| = \infty.$$

$$\sigma_p := \frac{1}{18} \text{tr } T_{\mathbf{k}}^{2p}, \quad F_{\mathbf{k}}(\alpha) := \det_2(I - \alpha^2 T_{\mathbf{k}}^2)$$

p	$\sqrt{3}\sigma_p/3^p\pi$
2	4/9
3	32/63
4	40/81

p	$\sqrt{3}\sigma_p/3^p\pi$
5	9560/20007
6	245120/527877
7	1957475168/4337177481

Theorem (BHZ '22) The largest real eigenvalue of $T_{\mathbf{k}}$, $1/\alpha_*$, is *simple* and $\alpha_* \in (0.583, 0.589)$.

$$\text{Spec}_{L^2(\mathbb{C}/\Gamma)} D(\alpha) = \begin{cases} \Gamma^* & \alpha \notin \mathcal{A} \\ \mathbb{C} & \alpha \in \mathcal{A}, \end{cases}$$

Theorem (BHZ '22) For all $p > 1$

$$\sum_{\alpha \in \mathcal{A}} \alpha^{-2p} \in \frac{\pi}{\sqrt{3}} \mathbb{Q} \quad \text{and as a consequence } |\mathcal{A}| = \infty.$$

$$\sigma_p := \frac{1}{18} \text{tr } T_{\mathbf{k}}^{2p}, \quad F_{\mathbf{k}}(\alpha) := \det_2(I - \alpha^2 T_{\mathbf{k}}^2)$$

p	$\sqrt{3}\sigma_p/3^p\pi$
2	4/9
3	32/63
4	40/81

p	$\sqrt{3}\sigma_p/3^p\pi$
5	9560/20007
6	245120/527877
7	1957475168/4337177481

Theorem (BHZ '22) The largest real eigenvalue of $T_{\mathbf{k}}$, $1/\alpha_*$, is *simple* and $\alpha_* \in (0.583, 0.589)$.

Consequently, the flat band of $H(\alpha_*)$ is *simple*.

Spectral characterization allows accurate computation of more α 's:

Spectral characterization allows accurate computation of more α 's:

k	α_k	$\alpha_k - \alpha_{k-1}$
1	0.58566355838955	
2	2.2211821738201	1.6355
3	3.7514055099052	1.5302
4	5.276497782985	1.5251
5	6.79478505720	1.5183
6	8.3129991933	1.5182
7	9.829066969	1.5161
8	11.34534068	1.5163
9	12.8606086	1.5153
10	14.376072	1.5155
11	15.89096	1.5149
12	17.4060	1.5150
13	18.920	1.5147

Spectral characterization allows accurate computation of more α 's:

k	α_k	$\alpha_k - \alpha_{k-1}$
1	0.58566355838955	
2	2.2211821738201	1.6355
3	3.7514055099052	1.5302
4	5.276497782985	1.5251
5	6.79478505720	1.5183
6	8.3129991933	1.5182
7	9.829066969	1.5161
8	11.34534068	1.5163
9	12.8606086	1.5153
10	14.376072	1.5155
11	15.89096	1.5149
12	17.4060	1.5150
13	18.920	1.5147

Numerically all of these are simple

Spectral characterization allows accurate computation of more α 's:


k	α_k	$\alpha_k - \alpha_{k-1}$
1	0.58566355838955	
2	2.2211821738201	1.6355
3	3.7514055099052	1.5302
4	5.276497782985	1.5251
5	6.79478505720	1.5183
6	8.3129991933	1.5182
7	9.829066969	1.5161
8	11.34534068	1.5163
9	12.8606086	1.5153
10	14.376072	1.5155
11	15.89096	1.5149
12	17.4060	1.5150
13	18.920	1.5147

Numerically all of these are **simple** That is **not** the case for complex α 's or for real α 's and more general potentials:

Spectral characterization allows accurate computation of more α 's:

k	α_k	$\alpha_k - \alpha_{k-1}$
1	0.58566355838955	
2	2.2211821738201	1.6355
3	3.7514055099052	1.5302
4	5.276497782985	1.5251
5	6.79478505720	1.5183
6	8.3129991933	1.5182
7	9.829066969	1.5161
8	11.34534068	1.5163
9	12.8606086	1.5153
10	14.376072	1.5155
11	15.89096	1.5149
12	17.4060	1.5150
13	18.920	1.5147

Numerically all of these are simple That is **not** the case for complex α 's or for real α 's and more general potentials:




Works for general potentials with $\mathbb{Z}_3^2 \rtimes \mathbb{Z}_3$ symmetries

$$U_\theta(z) := \sum_{k=0}^2 \omega^k (\cos^2 \theta e^{\frac{1}{2}(\bar{z}\omega^k - z\bar{\omega}^k)} + \sin^2 \theta e^{\bar{z}\omega^k - z\bar{\omega}^k})$$


Works for general potentials with $\mathbb{Z}_3^2 \times \mathbb{Z}_3$ symmetries

$$U_\theta(z) := \sum_{k=0}^2 \omega^k (\cos^2 \theta e^{\frac{1}{2}(\bar{z}\omega^k - z\bar{\omega}^k)} + \sin^2 \theta e^{\bar{z}\omega^k - z\bar{\omega}^k})$$



Works for general potentials with $\mathbb{Z}_3^2 \times \mathbb{Z}_3$ symmetries

$$U_\theta(z) := \sum_{k=0}^2 \omega^k (\cos^2 \theta e^{\frac{1}{2}(\bar{z}\omega^k - z\bar{\omega}^k)} + \sin^2 \theta e^{\bar{z}\omega^k - z\bar{\omega}^k})$$



Theorem (BHZ '22) *For a generic potential flat bands are simple.*

Flat bands from theta functions

Tarnopolsky et al '19: consider $\mathbf{u} \in L^2_{\rho_{1,0}}(\mathbb{C}/\Gamma; \mathbb{C}^2)$, $D(\alpha)\mathbf{u} = 0$

Flat bands from theta functions

Tarnopolsky et al '19: consider $\mathbf{u} \in L^2_{\rho_{1,0}}(\mathbb{C}/\Gamma; \mathbb{C}^2)$, $D(\alpha)\mathbf{u} = 0$

$\mathbf{u}_k(z) := e^{\frac{i}{2}(z\bar{k} + \bar{z}k)} f_k(z) \mathbf{u}(z)$, $z \mapsto e^{\frac{i}{2}(z\bar{k} + \bar{z}k)} f_k(z)$ periodic, $\partial_{\bar{z}} f_k = 0$

$$(D(\alpha) - \mathbf{k})\mathbf{u}_k(z) = 0$$

Problem: f_k with these properties will have poles

Flat bands from theta functions

Tarnopolsky et al '19: consider $\mathbf{u} \in L^2_{\rho_{1,0}}(\mathbb{C}/\Gamma; \mathbb{C}^2)$, $D(\alpha)\mathbf{u} = 0$

$\mathbf{u}_k(z) := e^{\frac{i}{2}(z\bar{k} + \bar{z}k)} f_k(z) \mathbf{u}(z)$, $z \mapsto e^{\frac{i}{2}(z\bar{k} + \bar{z}k)} f_k(z)$ periodic, $\partial_{\bar{z}} f_k = 0$

$$(D(\alpha) - \mathbf{k})\mathbf{u}_k(z) = 0$$

Problem: f_k with these properties will have poles

Solution: Look for α 's at which \mathbf{u} has a zero!


Flat bands from theta functions

Tarnopolsky et al '19: consider $\mathbf{u} \in L^2_{\rho_{1,0}}(\mathbb{C}/\Gamma; \mathbb{C}^2)$, $D(\alpha)\mathbf{u} = 0$

$$\mathbf{u}_k(z) := e^{\frac{i}{2}(z\bar{k} + \bar{z}k)} f_k(z) \mathbf{u}(z), \quad z \mapsto e^{\frac{i}{2}(z\bar{k} + \bar{z}k)} f_k(z) \text{ periodic, } \partial_{\bar{z}} f_k = 0$$
$$(D(\alpha) - \mathbf{k})\mathbf{u}_k(z) = 0$$

Problem: f_k with these properties will have poles

Solution: Look for α 's at which \mathbf{u} has a zero!



Flat bands from theta functions

Tarnopolsky et al '19: consider $\mathbf{u} \in L^2_{\rho_{1,0}}(\mathbb{C}/\Gamma; \mathbb{C}^2)$, $D(\alpha)\mathbf{u} = 0$

$\mathbf{u}_k(z) := e^{\frac{i}{2}(z\bar{k} + \bar{z}k)} f_k(z) \mathbf{u}(z)$, $z \mapsto e^{\frac{i}{2}(z\bar{k} + \bar{z}k)} f_k(z)$ periodic, $\partial_{\bar{z}} f_k = 0$

$$(D(\alpha) - \mathbf{k})\mathbf{u}_k(z) = 0$$

Problem: f_k with these properties will have poles

Solution: Look for α 's at which \mathbf{u} has a zero!

$$\mathbf{u}(\alpha, z_S) = 0, \quad \alpha \in \mathcal{A}, \quad z_S = \frac{4\sqrt{3}}{9}\pi, \quad z_S \equiv \omega z_S \pmod{\Gamma/3}$$

Flat bands from theta functions

Tarnopolsky et al '19: consider $\mathbf{u} \in L^2_{\rho_{1,0}}(\mathbb{C}/\Gamma; \mathbb{C}^2)$, $D(\alpha)\mathbf{u} = 0$

$\mathbf{u}_k(z) := e^{\frac{i}{2}(z\bar{k}+\bar{z}k)} f_k(z)\mathbf{u}(z)$, $z \mapsto e^{\frac{i}{2}(z\bar{k}+\bar{z}k)} f_k(z)$ periodic, $\partial_{\bar{z}} f_k = 0$

$$(D(\alpha) - \mathbf{k})\mathbf{u}_k(z) = 0$$

Problem: f_k with these properties will have poles

Solution: Look for α 's at which \mathbf{u} has a zero!

$$\mathbf{u}(\alpha, z_S) = 0, \quad \alpha \in \mathcal{A}, \quad z_S = \frac{4\sqrt{3}}{9}\pi, \quad z_S \equiv \omega z_S \pmod{\Gamma/3}$$

$$e^{\frac{i}{2}(z\bar{k}+\bar{z}k)} f_k(z) = e^{2\pi(\zeta-\bar{\zeta})k/\sqrt{3}} \frac{\theta_1(\zeta + k|\omega)}{\theta_1(\zeta|\omega)}, \quad z = \frac{4}{3}\pi i\omega\zeta$$

Flat bands from theta functions

Tarnopolsky et al '19: consider $\mathbf{u} \in L^2_{\rho_{1,0}}(\mathbb{C}/\Gamma; \mathbb{C}^2)$, $D(\alpha)\mathbf{u} = 0$

$\mathbf{u}_k(z) := e^{\frac{i}{2}(z\bar{k}+\bar{z}k)} f_k(z)\mathbf{u}(z)$, $z \mapsto e^{\frac{i}{2}(z\bar{k}+\bar{z}k)} f_k(z)$ periodic, $\partial_{\bar{z}} f_k = 0$

$$(D(\alpha) - \mathbf{k})\mathbf{u}_k(z) = 0$$

Problem: f_k with these properties will have poles

Solution: Look for α 's at which \mathbf{u} has a zero!

$$\mathbf{u}(\alpha, z_S) = 0, \quad \alpha \in \mathcal{A}, \quad z_S = \frac{4\sqrt{3}}{9}\pi, \quad z_S \equiv \omega z_S \pmod{\Gamma/3}$$

$$e^{\frac{i}{2}(z\bar{k}+\bar{z}k)} f_k(z) = e^{2\pi(\zeta-\bar{\zeta})k/\sqrt{3}} \frac{\theta_1(\zeta + k|\omega)}{\theta_1(\zeta|\omega)}, \quad z = \frac{4}{3}\pi i\omega\zeta$$

Similar argument in Dubrovin–Novikov '80

Flat bands from theta functions

Tarnopolsky et al '19: consider $\mathbf{u} \in L^2_{\rho_{1,0}}(\mathbb{C}/\Gamma; \mathbb{C}^2)$, $D(\alpha)\mathbf{u} = 0$

$\mathbf{u}_k(z) := e^{\frac{i}{2}(z\bar{k}+\bar{z}k)} f_k(z)\mathbf{u}(z)$, $z \mapsto e^{\frac{i}{2}(z\bar{k}+\bar{z}k)} f_k(z)$ periodic, $\partial_{\bar{z}} f_k = 0$

$$(D(\alpha) - \mathbf{k})\mathbf{u}_k(z) = 0$$

Problem: f_k with these properties will have poles

Solution: Look for α 's at which \mathbf{u} has a zero!

$$\mathbf{u}(\alpha, z_S) = 0, \quad \alpha \in \mathcal{A}, \quad z_S = \frac{4\sqrt{3}}{9}\pi, \quad z_S \equiv \omega z_S \pmod{\Gamma/3}$$

$$e^{\frac{i}{2}(z\bar{k}+\bar{z}k)} f_k(z) = e^{2\pi(\zeta-\bar{\zeta})k/\sqrt{3}} \frac{\theta_1(\zeta + k|\omega)}{\theta_1(\zeta|\omega)}, \quad z = \frac{4}{3}\pi i \omega \zeta$$

Similar argument in Dubrovin–Novikov '80

Theorem (BHZ '22) $\alpha \in \mathcal{A}$ simple $\Rightarrow z_S$ is the only zero of \mathbf{u} .

Flat bands from theta functions

$$e^{\frac{i}{2}(z\bar{\mathbf{k}}+\bar{z}\mathbf{k})} f_k(z) = e^{2\pi(\zeta-\bar{\zeta})k/\sqrt{3}} \frac{\theta_1(\zeta+k|\omega)}{\theta_1(\zeta|\omega)}, \quad z = \frac{4}{3}\pi i\omega\zeta$$

Flat bands from theta functions

$$e^{\frac{i}{2}(z\bar{k}+\bar{z}k)} f_k(z) = e^{2\pi(\zeta-\bar{\zeta})k/\sqrt{3}} \frac{\theta_1(\zeta+k|\omega)}{\theta_1(\zeta|\omega)}, \quad z = \frac{4}{3}\pi i\omega\zeta$$

$$\theta_1(\zeta|\omega) := - \sum_{n \in \mathbb{Z}} e^{\pi i(n+\frac{1}{2})^2\omega + 2\pi i(n+\frac{1}{2})(\zeta+\frac{1}{2})}$$

$$\theta_1(\zeta+m|\omega) = (-1)^m \theta_1(\zeta|\omega), \quad \theta_1(\zeta+n\omega|\omega) = (-1)^n e^{-\pi i n\omega - 2\pi i \zeta n} \theta_1(\zeta|\omega)$$

Flat bands from theta functions

$$e^{\frac{i}{2}(z\bar{k}+\bar{z}k)} f_k(z) = e^{2\pi(\zeta-\bar{\zeta})k/\sqrt{3}} \frac{\theta_1(\zeta+k|\omega)}{\theta_1(\zeta|\omega)}, \quad z = \frac{4}{3}\pi i\omega\zeta$$

$$\theta_1(\zeta|\omega) := - \sum_{n \in \mathbb{Z}} e^{\pi i(n+\frac{1}{2})^2\omega + 2\pi i(n+\frac{1}{2})(\zeta+\frac{1}{2})}$$

$$\theta_1(\zeta+m|\omega) = (-1)^m \theta_1(\zeta|\omega), \quad \theta_1(\zeta+n\omega|\omega) = (-1)^n e^{-\pi i n\omega - 2\pi i \zeta n} \theta_1(\zeta|\omega)$$

$$\theta_1(\zeta|\omega) = 0 \iff \zeta \in \mathbb{Z}\omega + \mathbb{Z}$$

Flat bands from theta functions

$$e^{\frac{i}{2}(z\bar{\mathbf{k}}+\bar{z}\mathbf{k})} f_{\mathbf{k}}(z) = e^{2\pi(\zeta-\bar{\zeta})k/\sqrt{3}} \frac{\theta_1(\zeta+k|\omega)}{\theta_1(\zeta|\omega)}, \quad z = \frac{4}{3}\pi i\omega\zeta$$

$$\theta_1(\zeta|\omega) := - \sum_{n \in \mathbb{Z}} e^{\pi i(n+\frac{1}{2})^2\omega + 2\pi i(n+\frac{1}{2})(\zeta+\frac{1}{2})}$$

$$\theta_1(\zeta+m|\omega) = (-1)^m \theta_1(\zeta|\omega), \quad \theta_1(\zeta+n\omega|\omega) = (-1)^n e^{-\pi i n\omega - 2\pi i \zeta n} \theta_1(\zeta|\omega)$$

$$\theta_1(\zeta|\omega) = 0 \iff \zeta \in \mathbb{Z}\omega + \mathbb{Z}$$

$\mathbf{k} \mapsto u_{\mathbf{k}} = e^{\frac{i}{2}(z\bar{\mathbf{k}}+\bar{z}\mathbf{k})} f_{\mathbf{k}}(z) \mathbf{u}(z)$ is holomorphic

Flat bands from theta functions

$$e^{\frac{i}{2}(z\bar{\mathbf{k}}+\bar{z}\mathbf{k})} f_{\mathbf{k}}(z) = e^{2\pi(\zeta-\bar{\zeta})\mathbf{k}/\sqrt{3}} \frac{\theta_1(\zeta + \mathbf{k}|\omega)}{\theta_1(\zeta|\omega)}, \quad z = \frac{4}{3}\pi i\omega\zeta$$

$$\theta_1(\zeta|\omega) := - \sum_{n \in \mathbb{Z}} e^{\pi i(n+\frac{1}{2})^2\omega + 2\pi i(n+\frac{1}{2})(\zeta+\frac{1}{2})}$$

$$\theta_1(\zeta+m|\omega) = (-1)^m \theta_1(\zeta|\omega), \quad \theta_1(\zeta+n\omega|\omega) = (-1)^n e^{-\pi i n\omega - 2\pi i \zeta n} \theta_1(\zeta|\omega)$$

$$\theta_1(\zeta|\omega) = 0 \iff \zeta \in \mathbb{Z}\omega + \mathbb{Z}$$

$\mathbf{k} \mapsto u_{\mathbf{k}} = e^{\frac{i}{2}(z\bar{\mathbf{k}}+\bar{z}\mathbf{k})} f_{\mathbf{k}}(z) \mathbf{u}(z)$ is holomorphic **Ledwith et al '21**

Flat bands from theta functions

$$e^{\frac{i}{2}(z\bar{\mathbf{k}}+\bar{z}\mathbf{k})} f_k(z) = e^{2\pi(\zeta-\bar{\zeta})k/\sqrt{3}} \frac{\theta_1(\zeta+k|\omega)}{\theta_1(\zeta|\omega)}, \quad z = \frac{4}{3}\pi i\omega\zeta$$

$$\theta_1(\zeta|\omega) := - \sum_{n \in \mathbb{Z}} e^{\pi i(n+\frac{1}{2})^2\omega+2\pi i(n+\frac{1}{2})(\zeta+\frac{1}{2})}$$

$$\theta_1(\zeta+m|\omega) = (-1)^m \theta_1(\zeta|\omega), \quad \theta_1(\zeta+n\omega|\omega) = (-1)^n e^{-\pi i n\omega-2\pi i\zeta n} \theta_1(\zeta|\omega)$$

$$\theta_1(\zeta|\omega) = 0 \iff \zeta \in \mathbb{Z}\omega + \mathbb{Z}$$

$\mathbf{k} \mapsto u_{\mathbf{k}} = e^{\frac{i}{2}(z\bar{\mathbf{k}}+\bar{z}\mathbf{k})} f_k(z) \mathbf{u}(z)$ is holomorphic **Ledwith et al '21**

Felix Klein 1920: *When I was a student, abelian functions were, as an effect of the Jacobian tradition, considered the uncontested summit of mathematics, and each of us was ambitious to make progress in this field. And now? The younger generation hardly knows abelian functions.*


Fine structure of eigenfunctions

Fine structure of eigenfunctions

For $\mathbf{k} \in \Gamma^*/3\Gamma^*$ we determine which representation of G_3 (Heisenberg group over \mathbb{Z}_3), $u_{\mathbf{k}}$ falls into:

Fine structure of eigenfunctions


For $\mathbf{k} \in \Gamma^*/3\Gamma^*$ we determine which representation of G_3 (Heisenberg group over \mathbb{Z}_3), $u_{\mathbf{k}}$ falls into:




$\log |w|$ of the two component of $\mathbf{u}_0 \in L^2_{\rho_{0,0}} \subset L^2_0$, both vanishing at $-z_5$. Except for vanishing at $-z_5$ this eigenstate exists for all α 's.

Fine structure of eigenfunctions

For $\mathbf{k} \in \Gamma^*/3\Gamma^*$ we determine which representation of G_3 (Heisenberg group over \mathbb{Z}_3), $u_{\mathbf{k}}$ falls into:




$\log |w|$ of the two component of $\mathbf{u}_0 \in L^2_{\rho_{0,0}} \subset L^2_0$, both vanishing at $-z_5$. Except for vanishing at $-z_5$ this eigenstate exists for all α 's.




$\log |w|$ of components of $\mathbf{u}_i \in L^2_{\rho_{2,1}} \subset L^2_i$.

Fine structure of eigenfunctions

For $\mathbf{k} \in \Gamma^*/3\Gamma^*$ we determine which representation of G_3 (Heisenberg group over \mathbb{Z}_3), $u_{\mathbf{k}}$ falls into:



$\log |w|$ of the two component of $\mathbf{u}_0 \in L^2_{\rho_{0,0}} \subset L^2_0$, both vanishing at $-z_5$. Except for vanishing at $-z_5$ this eigenstate exists for all α 's.




$\log |w|$ of components of $\mathbf{u}_i \in L^2_{\rho_{2,1}} \subset L^2_i$.

This is the state considered by [Cano et al '21](#) with additional symmetry $[\psi(z), \varphi(z)] \mapsto [-\varphi(-z), \psi(-z)]$.

Fine structure of eigenfunctions

Fine structure of eigenfunctions


Three dimensional representations come from $\{\pm\omega^k/\sqrt{3}\}_{k=0}^2$:



$\alpha = \alpha_1$ (first magic angle): vanishing determines symmetries

Fine structure of eigenfunctions

Three dimensional representations come from $\{\pm\omega^k/\sqrt{3}\}_{k=0}^2$:



$\alpha = \alpha_5$ (fifth magic angle): vanishing determines symmetries;
despite appearances $\mathbf{u}_{\omega^k/\sqrt{3}}$ do not vanish at other points.

Chern connection and curvature

$\mathbb{C}/3\Gamma^* \ni \mathbf{k} \rightarrow u_{\mathbf{k}} \in L^2_0(\mathbb{C}/\Gamma)$ defines a holomorphic line bundle

Ledwith et al '21

Chern connection and curvature

$\mathbb{C}/3\Gamma^* \ni \mathbf{k} \rightarrow u_{\mathbf{k}} \in L^2_0(\mathbb{C}/\Gamma)$ defines a holomorphic line bundle

Ledwith et al '21

Chern connection: $\eta := \partial_{\mathbf{k}} \log \|u_{\mathbf{k}}\|^2$

Chern connection and curvature

$\mathbb{C}/3\Gamma^* \ni \mathbf{k} \rightarrow u_{\mathbf{k}} \in L^2_0(\mathbb{C}/\Gamma)$ defines a holomorphic line bundle

Ledwith et al '21

Chern connection: $\eta := \partial_{\mathbf{k}} \log \|u_{\mathbf{k}}\|^2 = \|\mathbf{u}_{\mathbf{k}}\|^{-2} \langle \partial_{\mathbf{k}} \mathbf{u}_{\mathbf{k}}, \mathbf{u}_{\mathbf{k}} \rangle d\mathbf{k}$

Chern connection and curvature

$\mathbb{C}/3\Gamma^* \ni \mathbf{k} \rightarrow u_{\mathbf{k}} \in L^2_0(\mathbb{C}/\Gamma)$ defines a holomorphic line bundle

Ledwith et al '21

Chern connection: $\eta := \partial_{\mathbf{k}} \log \|u_{\mathbf{k}}\|^2 = \|u_{\mathbf{k}}\|^{-2} \langle \partial_{\mathbf{k}} u_{\mathbf{k}}, u_{\mathbf{k}} \rangle d\mathbf{k}$

Curvature: $\Omega = d\eta = \bar{\partial}_{\mathbf{k}} \partial_{\mathbf{k}} \log \|u_{\mathbf{k}}\|^2$

Chern connection and curvature

$\mathbb{C}/3\Gamma^* \ni \mathbf{k} \rightarrow u_{\mathbf{k}} \in L^2_0(\mathbb{C}/\Gamma)$ defines a holomorphic line bundle

Ledwith et al '21

Chern connection: $\eta := \partial_{\mathbf{k}} \log \|u_{\mathbf{k}}\|^2 = \|\mathbf{u}_{\mathbf{k}}\|^{-2} \langle \partial_{\mathbf{k}} \mathbf{u}_{\mathbf{k}}, \mathbf{u}_{\mathbf{k}} \rangle d\mathbf{k}$

Curvature: $\Omega = d\eta = \bar{\partial}_{\mathbf{k}} \partial_{\mathbf{k}} \log \|u_{\mathbf{k}}\|^2 = H(\mathbf{k}) d\bar{\mathbf{k}} \wedge d\mathbf{k}, \quad H(\mathbf{k}) \geq 0.$

Chern connection and curvature

$\mathbb{C}/3\Gamma^* \ni \mathbf{k} \rightarrow u_{\mathbf{k}} \in L^2_0(\mathbb{C}/\Gamma)$ defines a holomorphic line bundle

Ledwith et al '21

Chern connection: $\eta := \partial_{\mathbf{k}} \log \|u_{\mathbf{k}}\|^2 = \|\mathbf{u}_{\mathbf{k}}\|^{-2} \langle \partial_{\mathbf{k}} \mathbf{u}_{\mathbf{k}}, \mathbf{u}_{\mathbf{k}} \rangle d\mathbf{k}$

Curvature: $\Omega = d\eta = \bar{\partial}_{\mathbf{k}} \partial_{\mathbf{k}} \log \|u_{\mathbf{k}}\|^2 = H(\mathbf{k}) d\bar{\mathbf{k}} \wedge d\mathbf{k}$, $H(\mathbf{k}) \geq 0$.

Chern class: $c_1 = \frac{i}{2\pi} \int_{\mathbb{C}/3\Gamma^*} \Omega$

Chern connection and curvature

$\mathbb{C}/3\Gamma^* \ni \mathbf{k} \rightarrow u_{\mathbf{k}} \in L^2_0(\mathbb{C}/\Gamma)$ defines a holomorphic line bundle

Ledwith et al '21

Chern connection: $\eta := \partial_{\mathbf{k}} \log \|u_{\mathbf{k}}\|^2 = \|u_{\mathbf{k}}\|^{-2} \langle \partial_{\mathbf{k}} u_{\mathbf{k}}, u_{\mathbf{k}} \rangle d\mathbf{k}$

Curvature: $\Omega = d\eta = \bar{\partial}_{\mathbf{k}} \partial_{\mathbf{k}} \log \|u_{\mathbf{k}}\|^2 = H(\mathbf{k}) d\bar{\mathbf{k}} \wedge d\mathbf{k}, \quad H(\mathbf{k}) \geq 0.$

Chern class: $c_1 = \frac{i}{2\pi} \int_{\mathbb{C}/3\Gamma^*} \Omega = -1$

Chern connection and curvature

$\mathbb{C}/3\Gamma^* \ni \mathbf{k} \rightarrow u_{\mathbf{k}} \in L^2_0(\mathbb{C}/\Gamma)$ defines a holomorphic line bundle

Ledwith et al '21

Chern connection: $\eta := \partial_{\mathbf{k}} \log \|u_{\mathbf{k}}\|^2 = \|\mathbf{u}_{\mathbf{k}}\|^{-2} \langle \partial_{\mathbf{k}} \mathbf{u}_{\mathbf{k}}, \mathbf{u}_{\mathbf{k}} \rangle d\mathbf{k}$

Curvature: $\Omega = d\eta = \bar{\partial}_{\mathbf{k}} \partial_{\mathbf{k}} \log \|u_{\mathbf{k}}\|^2 = H(\mathbf{k}) d\bar{\mathbf{k}} \wedge d\mathbf{k}, \quad H(\mathbf{k}) \geq 0.$

Chern class: $c_1 = \frac{i}{2\pi} \int_{\mathbb{C}/3\Gamma^*} \Omega = -1$

Standard deviation: $|\mathbb{C}/3\Gamma^*|^{-1} \int_{\mathbb{C}/3\Gamma^*} (H(k)/2\pi + c_1)^2 dm(\mathbf{k})$

Chern connection and curvature

$\mathbb{C}/3\Gamma^* \ni \mathbf{k} \rightarrow u_{\mathbf{k}} \in L^2_0(\mathbb{C}/\Gamma)$ defines a holomorphic line bundle

Ledwith et al '21

Chern connection: $\eta := \partial_{\mathbf{k}} \log \|u_{\mathbf{k}}\|^2 = \|u_{\mathbf{k}}\|^{-2} \langle \partial_{\mathbf{k}} u_{\mathbf{k}}, u_{\mathbf{k}} \rangle d\mathbf{k}$

Curvature: $\Omega = d\eta = \bar{\partial}_{\mathbf{k}} \partial_{\mathbf{k}} \log \|u_{\mathbf{k}}\|^2 = H(\mathbf{k}) d\bar{\mathbf{k}} \wedge d\mathbf{k}$, $H(\mathbf{k}) \geq 0$.

Chern class: $c_1 = \frac{i}{2\pi} \int_{\mathbb{C}/3\Gamma^*} \Omega = -1$

Standard deviation: $|\mathbb{C}/3\Gamma^*|^{-1} \int_{\mathbb{C}/3\Gamma^*} (H(k)/2\pi + c_1)^2 dm(\mathbf{k})$

<i>Magic angle</i>	0.58	2.22	3.75	5.27	6.79	8.31
<i>std</i>	0.2834	0.1502	0.1333	0.1611	0.1965	0.2362
<i>Magic angle</i>	9.82	11.34	12.86	14.37	15.89	17.40
<i>std</i>	0.2674	0.2904	0.3518	0.3835	0.3346	0.2790

Chern connection and curvature

$\mathbb{C}/3\Gamma^* \ni \mathbf{k} \rightarrow u_{\mathbf{k}} \in L_0^2(\mathbb{C}/\Gamma)$ defines a holomorphic line bundle


Ledwith et al '21

Chern connection: $\eta := \partial_{\mathbf{k}} \log \|u_{\mathbf{k}}\|^2 = \|u_{\mathbf{k}}\|^{-2} \langle \partial_{\mathbf{k}} u_{\mathbf{k}}, u_{\mathbf{k}} \rangle d\mathbf{k}$

Curvature: $\Omega = d\eta = \bar{\partial}_{\mathbf{k}} \partial_{\mathbf{k}} \log \|u_{\mathbf{k}}\|^2 = H(\mathbf{k}) d\bar{\mathbf{k}} \wedge d\mathbf{k}, \quad H(\mathbf{k}) \geq 0.$

Chern class: $c_1 = \frac{i}{2\pi} \int_{\mathbb{C}/3\Gamma^*} \Omega = -1$

Standard deviation: $|\mathbb{C}/3\Gamma^*|^{-1} \int_{\mathbb{C}/3\Gamma^*} (H(k)/2\pi + c_1)^2 dm(\mathbf{k})$



$\alpha_1 \simeq 0.58$; concentration at \mathbf{i}

Chern connection and curvature

$\mathbb{C}/3\Gamma^* \ni \mathbf{k} \rightarrow u_{\mathbf{k}} \in L_0^2(\mathbb{C}/\Gamma)$ defines a holomorphic line bundle


Ledwith et al '21

Chern connection: $\eta := \partial_{\mathbf{k}} \log \|u_{\mathbf{k}}\|^2 = \|u_{\mathbf{k}}\|^{-2} \langle \partial_{\mathbf{k}} u_{\mathbf{k}}, u_{\mathbf{k}} \rangle d\mathbf{k}$

Curvature: $\Omega = d\eta = \bar{\partial}_{\mathbf{k}} \partial_{\mathbf{k}} \log \|u_{\mathbf{k}}\|^2 = H(\mathbf{k}) d\bar{\mathbf{k}} \wedge d\mathbf{k}, \quad H(\mathbf{k}) \geq 0.$

Chern class: $c_1 = \frac{i}{2\pi} \int_{\mathbb{C}/3\Gamma^*} \Omega = -1$

Standard deviation: $|\mathbb{C}/3\Gamma^*|^{-1} \int_{\mathbb{C}/3\Gamma^*} (H(k)/2\pi + c_1)^2 dm(\mathbf{k})$



$\alpha_1 \simeq 0.58$; concentration at $\mathbf{i} \equiv \omega i \pmod{3\Gamma^*}$

Chern connection and curvature

$\mathbb{C}/3\Gamma^* \ni \mathbf{k} \rightarrow u_{\mathbf{k}} \in L_0^2(\mathbb{C}/\Gamma)$ defines a holomorphic line bundle


Ledwith et al '21

Chern connection: $\eta := \partial_{\mathbf{k}} \log \|u_{\mathbf{k}}\|^2 = \|u_{\mathbf{k}}\|^{-2} \langle \partial_{\mathbf{k}} u_{\mathbf{k}}, u_{\mathbf{k}} \rangle d\mathbf{k}$

Curvature: $\Omega = d\eta = \bar{\partial}_{\mathbf{k}} \partial_{\mathbf{k}} \log \|u_{\mathbf{k}}\|^2 = H(\mathbf{k}) d\bar{\mathbf{k}} \wedge d\mathbf{k}, \quad H(\mathbf{k}) \geq 0.$

Chern class: $c_1 = \frac{i}{2\pi} \int_{\mathbb{C}/3\Gamma^*} \Omega = -1$

Standard deviation: $|\mathbb{C}/3\Gamma^*|^{-1} \int_{\mathbb{C}/3\Gamma^*} (H(k)/2\pi + c_1)^2 dm(\mathbf{k})$



$\alpha_2 \simeq 2.22$; concentration at $\mathbf{i} \equiv \omega i \pmod{3\Gamma^*}$

Chern connection and curvature

$\mathbb{C}/3\Gamma^* \ni \mathbf{k} \rightarrow u_{\mathbf{k}} \in L_0^2(\mathbb{C}/\Gamma)$ defines a holomorphic line bundle


Ledwith et al '21

Chern connection: $\eta := \partial_{\mathbf{k}} \log \|u_{\mathbf{k}}\|^2 = \|u_{\mathbf{k}}\|^{-2} \langle \partial_{\mathbf{k}} u_{\mathbf{k}}, u_{\mathbf{k}} \rangle d\mathbf{k}$

Curvature: $\Omega = d\eta = \bar{\partial}_{\mathbf{k}} \partial_{\mathbf{k}} \log \|u_{\mathbf{k}}\|^2 = H(\mathbf{k}) d\bar{\mathbf{k}} \wedge d\mathbf{k}, \quad H(\mathbf{k}) \geq 0.$

Chern class: $c_1 = \frac{i}{2\pi} \int_{\mathbb{C}/3\Gamma^*} \Omega = -1$

Standard deviation: $|\mathbb{C}/3\Gamma^*|^{-1} \int_{\mathbb{C}/3\Gamma^*} (H(k)/2\pi + c_1)^2 dm(\mathbf{k})$



$\alpha_3 \simeq 3.75$; concentration at $\mathbf{i} \equiv \omega i \pmod{3\Gamma^*}$

Chern connection and curvature

$\mathbb{C}/3\Gamma^* \ni \mathbf{k} \rightarrow u_{\mathbf{k}} \in L_0^2(\mathbb{C}/\Gamma)$ defines a holomorphic line bundle


Ledwith et al '21

Chern connection: $\eta := \partial_{\mathbf{k}} \log \|u_{\mathbf{k}}\|^2 = \|u_{\mathbf{k}}\|^{-2} \langle \partial_{\mathbf{k}} u_{\mathbf{k}}, u_{\mathbf{k}} \rangle d\mathbf{k}$

Curvature: $\Omega = d\eta = \bar{\partial}_{\mathbf{k}} \partial_{\mathbf{k}} \log \|u_{\mathbf{k}}\|^2 = H(\mathbf{k}) d\bar{\mathbf{k}} \wedge d\mathbf{k}, \quad H(\mathbf{k}) \geq 0.$

Chern class: $c_1 = \frac{i}{2\pi} \int_{\mathbb{C}/3\Gamma^*} \Omega = -1$

Standard deviation: $|\mathbb{C}/3\Gamma^*|^{-1} \int_{\mathbb{C}/3\Gamma^*} (H(k)/2\pi + c_1)^2 dm(\mathbf{k})$



$\alpha_4 \simeq 5.27$; concentration at $\mathbf{i} \equiv \omega i \pmod{3\Gamma^*}$

Chern connection and curvature

$\mathbb{C}/3\Gamma^* \ni \mathbf{k} \rightarrow u_{\mathbf{k}} \in L_0^2(\mathbb{C}/\Gamma)$ defines a holomorphic line bundle


Ledwith et al '21

Chern connection: $\eta := \partial_{\mathbf{k}} \log \|u_{\mathbf{k}}\|^2 = \|u_{\mathbf{k}}\|^{-2} \langle \partial_{\mathbf{k}} u_{\mathbf{k}}, u_{\mathbf{k}} \rangle d\mathbf{k}$

Curvature: $\Omega = d\eta = \bar{\partial}_{\mathbf{k}} \partial_{\mathbf{k}} \log \|u_{\mathbf{k}}\|^2 = H(\mathbf{k}) d\bar{\mathbf{k}} \wedge d\mathbf{k}$, $H(\mathbf{k}) \geq 0$.

Chern class: $c_1 = \frac{i}{2\pi} \int_{\mathbb{C}/3\Gamma^*} \Omega = -1$

Standard deviation: $|\mathbb{C}/3\Gamma^*|^{-1} \int_{\mathbb{C}/3\Gamma^*} (H(k)/2\pi + c_1)^2 dm(\mathbf{k})$



$\alpha_5 \simeq 6.79$; concentration at $\mathbf{i} \equiv \omega i \pmod{3\Gamma^*}$

Chern connection and curvature

$\mathbb{C}/3\Gamma^* \ni \mathbf{k} \rightarrow u_{\mathbf{k}} \in L_0^2(\mathbb{C}/\Gamma)$ defines a holomorphic line bundle


Ledwith et al '21

Chern connection: $\eta := \partial_{\mathbf{k}} \log \|u_{\mathbf{k}}\|^2 = \|u_{\mathbf{k}}\|^{-2} \langle \partial_{\mathbf{k}} u_{\mathbf{k}}, u_{\mathbf{k}} \rangle d\mathbf{k}$

Curvature: $\Omega = d\eta = \bar{\partial}_{\mathbf{k}} \partial_{\mathbf{k}} \log \|u_{\mathbf{k}}\|^2 = H(\mathbf{k}) d\bar{\mathbf{k}} \wedge d\mathbf{k}, \quad H(\mathbf{k}) \geq 0.$

Chern class: $c_1 = \frac{i}{2\pi} \int_{\mathbb{C}/3\Gamma^*} \Omega = -1$

Standard deviation: $|\mathbb{C}/3\Gamma^*|^{-1} \int_{\mathbb{C}/3\Gamma^*} (H(k)/2\pi + c_1)^2 dm(\mathbf{k})$



$\alpha_6 \simeq 8.31$; concentration at $\mathbf{i} \equiv \omega i \pmod{3\Gamma^*}$

Chern connection and curvature

$\mathbb{C}/3\Gamma^* \ni \mathbf{k} \rightarrow u_{\mathbf{k}} \in L_0^2(\mathbb{C}/\Gamma)$ defines a holomorphic line bundle


Ledwith et al '21

Chern connection: $\eta := \partial_{\mathbf{k}} \log \|u_{\mathbf{k}}\|^2 = \|u_{\mathbf{k}}\|^{-2} \langle \partial_{\mathbf{k}} u_{\mathbf{k}}, u_{\mathbf{k}} \rangle d\mathbf{k}$

Curvature: $\Omega = d\eta = \bar{\partial}_{\mathbf{k}} \partial_{\mathbf{k}} \log \|u_{\mathbf{k}}\|^2 = H(\mathbf{k}) d\bar{\mathbf{k}} \wedge d\mathbf{k}, \quad H(\mathbf{k}) \geq 0.$

Chern class: $c_1 = \frac{i}{2\pi} \int_{\mathbb{C}/3\Gamma^*} \Omega = -1$

Standard deviation: $|\mathbb{C}/3\Gamma^*|^{-1} \int_{\mathbb{C}/3\Gamma^*} (H(k)/2\pi + c_1)^2 dm(\mathbf{k})$



$\alpha_7 \simeq 9.82$; concentration at $\mathbf{i} \equiv \omega i \pmod{3\Gamma^*}$

Chern connection and curvature

$\mathbb{C}/3\Gamma^* \ni \mathbf{k} \rightarrow u_{\mathbf{k}} \in L_0^2(\mathbb{C}/\Gamma)$ defines a holomorphic line bundle


Ledwith et al '21

Chern connection: $\eta := \partial_{\mathbf{k}} \log \|u_{\mathbf{k}}\|^2 = \|u_{\mathbf{k}}\|^{-2} \langle \partial_{\mathbf{k}} u_{\mathbf{k}}, u_{\mathbf{k}} \rangle d\mathbf{k}$

Curvature: $\Omega = d\eta = \bar{\partial}_{\mathbf{k}} \partial_{\mathbf{k}} \log \|u_{\mathbf{k}}\|^2 = H(\mathbf{k}) d\bar{\mathbf{k}} \wedge d\mathbf{k}, \quad H(\mathbf{k}) \geq 0.$

Chern class: $c_1 = \frac{i}{2\pi} \int_{\mathbb{C}/3\Gamma^*} \Omega = -1$

Standard deviation: $|\mathbb{C}/3\Gamma^*|^{-1} \int_{\mathbb{C}/3\Gamma^*} (H(k)/2\pi + c_1)^2 dm(\mathbf{k})$



$\alpha_8 \simeq 11.34$; concentration at $\mathbf{i} \equiv \omega \mathbf{i} \pmod{3\Gamma^*}$

Chern connection and curvature

$\mathbb{C}/3\Gamma^* \ni \mathbf{k} \rightarrow u_{\mathbf{k}} \in L_0^2(\mathbb{C}/\Gamma)$ defines a holomorphic line bundle


Ledwith et al '21

Chern connection: $\eta := \partial_{\mathbf{k}} \log \|u_{\mathbf{k}}\|^2 = \|u_{\mathbf{k}}\|^{-2} \langle \partial_{\mathbf{k}} u_{\mathbf{k}}, u_{\mathbf{k}} \rangle d\mathbf{k}$

Curvature: $\Omega = d\eta = \bar{\partial}_{\mathbf{k}} \partial_{\mathbf{k}} \log \|u_{\mathbf{k}}\|^2 = H(\mathbf{k}) d\bar{\mathbf{k}} \wedge d\mathbf{k}, \quad H(\mathbf{k}) \geq 0.$

Chern class: $c_1 = \frac{i}{2\pi} \int_{\mathbb{C}/3\Gamma^*} \Omega = -1$

Standard deviation: $|\mathbb{C}/3\Gamma^*|^{-1} \int_{\mathbb{C}/3\Gamma^*} (H(k)/2\pi + c_1)^2 dm(\mathbf{k})$



$\alpha_g \simeq 12.96$; concentration at $\mathbf{i} \equiv \omega i \pmod{3\Gamma^*}$

Chern connection and curvature

$\mathbb{C}/3\Gamma^* \ni \mathbf{k} \rightarrow u_{\mathbf{k}} \in L_0^2(\mathbb{C}/\Gamma)$ defines a holomorphic line bundle


Ledwith et al '21

Chern connection: $\eta := \partial_{\mathbf{k}} \log \|u_{\mathbf{k}}\|^2 = \|u_{\mathbf{k}}\|^{-2} \langle \partial_{\mathbf{k}} u_{\mathbf{k}}, u_{\mathbf{k}} \rangle d\mathbf{k}$

Curvature: $\Omega = d\eta = \bar{\partial}_{\mathbf{k}} \partial_{\mathbf{k}} \log \|u_{\mathbf{k}}\|^2 = H(\mathbf{k}) d\bar{\mathbf{k}} \wedge d\mathbf{k}, \quad H(\mathbf{k}) \geq 0.$

Chern class: $c_1 = \frac{i}{2\pi} \int_{\mathbb{C}/3\Gamma^*} \Omega = -1$

Standard deviation: $|\mathbb{C}/3\Gamma^*|^{-1} \int_{\mathbb{C}/3\Gamma^*} (H(k)/2\pi + c_1)^2 dm(\mathbf{k})$




$\alpha_{10} \simeq 14.37$; concentration at $\mathbf{i} \equiv \omega i \pmod{3\Gamma^*}$

Some mathematical open problems

- ▶ Multiplicity issues; a stronger generic simplicity statement


Some mathematical open problems

- ▶ Multiplicity issues; a stronger generic simplicity statement



Some mathematical open problems


- ▶ Multiplicity issues; a stronger generic simplicity statement



- ▶ The fixed “shape” of the first band; what is a heuristic explanation?

Some mathematical open problems


- ▶ Multiplicity issues; a stronger generic simplicity statement



- ▶ The fixed “shape” of the first band; what is a heuristic explanation?
- ▶ Significance and explanation of the curvature “peak” at $\mathbf{k} = \mathbf{i}$

Some mathematical open problems


- ▶ Multiplicity issues; a stronger generic simplicity statement



- ▶ The fixed “shape” of the first band; what is a heuristic explanation?
- ▶ Significance and explanation of the curvature “peak” at $\mathbf{k} = \mathbf{i}$
- ▶ Asymptotics of $\alpha \in \mathcal{A} \cap \mathbb{R}_+$; in particular $\Delta\alpha \simeq \frac{3}{2}$?

Some mathematical open problems


- ▶ Multiplicity issues; a stronger generic simplicity statement



- ▶ The fixed “shape” of the first band; what is a heuristic explanation?
- ▶ Significance and explanation of the curvature “peak” at $\mathbf{k} = \mathbf{i}$
- ▶ Asymptotics of $\alpha \in \mathcal{A} \cap \mathbb{R}_+$; in particular $\Delta\alpha \simeq \frac{3}{2}$? Help from **Hitrik–Sjöstrand** '04... '?

Some mathematical open problems

- ▶ Multiplicity issues; a stronger generic simplicity statement



- ▶ The fixed “shape” of the first band; what is a heuristic explanation?
- ▶ Significance and explanation of the curvature “peak” at $\mathbf{k} = \mathbf{i}$
- ▶ Asymptotics of $\alpha \in \mathcal{A} \cap \mathbb{R}_+$; in particular $\Delta\alpha \simeq \frac{3}{2}$? Help from **Hitrik–Sjöstrand** '04... '?

