

The ground state energy of the dilute Bose gas.

Søren Fournais

Department of Mathematics, Aarhus University, Denmark
Based on joint work with Jan Philip Solovej, Copenhagen University, Denmark,
and Théotime Girardot, Lukas Junge, Léo Morin and Marco Olivieri, Aarhus.

Mathematical results of many-body quantum systems,
Herrsching 10/6 2022



AARHUS UNIVERSITY

Søren Fournais



The dilute Bose gas

Consider N interacting, non-relativistic bosons in a box Λ of sidelength L .

Density $\rho := N/|\Lambda| = N/L^d$, $d = 2, 3$.

The Hamiltonian of the system is, on the symmetric (bosonic) space $\otimes_s^N L^2(\Lambda)$,

$$H_N := \sum_{i=1}^N -\Delta_i + \sum_{i < j} v(x_i - x_j),$$

and $0 \leq v$ is radially symmetric with compact support.

The dilute Bose gas

Consider N interacting, non-relativistic bosons in a box Λ of sidelength L .

Density $\rho := N/|\Lambda| = N/L^d$, $d = 2, 3$.

The Hamiltonian of the system is, on the symmetric (bosonic) space $\otimes_s^N L^2(\Lambda)$,

$$H_N := \sum_{i=1}^N -\Delta_i + \sum_{i < j} v(x_i - x_j),$$

and $0 \leq v$ is radially symmetric with compact support.

The ground state energy of the system is

$$E_0(N, \Lambda) := \inf \text{Spec } H_N.$$

The dilute Bose gas

Consider N interacting, non-relativistic bosons in a box Λ of sidelength L .

Density $\rho := N/|\Lambda| = N/L^d$, $d = 2, 3$.

The Hamiltonian of the system is, on the symmetric (bosonic) space $\otimes_s^N L^2(\Lambda)$,

$$H_N := \sum_{i=1}^N -\Delta_i + \sum_{i < j} v(x_i - x_j),$$

and $0 \leq v$ is radially symmetric with compact support.

The ground state energy of the system is

$$E_0(N, \Lambda) := \inf \text{Spec } H_N.$$

The energy density in the thermodynamic limit is

$$e(\rho) = \lim_{L \rightarrow \infty, N/|\Lambda| = \rho} E_0(N, \Lambda)/L^d.$$

The scattering length l

Potential v is radial, positive with compact support.

The scattering length l

Potential v is radial, positive with compact support.

$$-\Delta\varphi + \frac{1}{2}v\varphi = 0,$$

where

$$\varphi = \begin{cases} 1 - a/r, & d = 3, \\ \log(r/a), & d = 2, \end{cases} \quad \text{outside } \text{supp } v.$$

Example:

$$v_{\text{hc}}(x) = \begin{cases} +\infty, & |x| < a, \\ 0, & |x| > a. \end{cases}$$

Here $\varphi = 0$ inside hard core and the scattering length equals the radius of the core.

The scattering length II

Let $\varphi_\delta = 2\delta\varphi$, with $\delta = \frac{1}{2}$ ($d = 3$) and $\delta \approx \frac{1}{|\log(\rho a^2)|}$ ($d = 2$) and $\omega = 1 - \varphi_\delta$.

Then, with $g = v(1 - \omega) = v\varphi_\delta$, the scattering equation can be reformulated as

$$-\Delta\omega = \frac{1}{2}g, \quad \text{i.e.} \quad \widehat{\omega}(k) = \frac{\widehat{g}(k)}{2k^2}, \quad k \neq 0,$$

and

$$8\pi a = \int_{\mathbb{R}^3} g < \int_{\mathbb{R}^3} v, \quad d = 3,$$

$$8\pi\delta = \int_{\mathbb{R}^2} g \ll \int_{\mathbb{R}^2} v, \quad d = 2.$$



The 3D Lee-Huang-Yang formula

Theorem

In the dilute limit $\rho a^3 \rightarrow 0$,

$$e^{3D}(\rho) = 4\pi\rho^2 a \left(1 + \frac{128}{15\sqrt{\pi}} \sqrt{\rho a^3} \right) + o(\rho^2 a (\rho a^3)^{1/2}).$$

- Lenz (1929), Bogoliubov (1947), Lee-Huang-Yang (1957).

The 3D Lee-Huang-Yang formula

Theorem

In the dilute limit $\rho a^3 \rightarrow 0$,

$$e^{3D}(\rho) = 4\pi\rho^2 a \left(1 + \frac{128}{15\sqrt{\pi}} \sqrt{\rho a^3} \right) + o(\rho^2 a (\rho a^3)^{1/2}).$$

- Lenz (1929), Bogoliubov (1947), Lee-Huang-Yang (1957).
- Rigorous proof of leading term Dyson (1957, upper), Lieb-Yngvason (1998).

The 3D Lee-Huang-Yang formula

Theorem

In the dilute limit $\rho a^3 \rightarrow 0$,

$$e^{3D}(\rho) = 4\pi\rho^2 a \left(1 + \frac{128}{15\sqrt{\pi}} \sqrt{\rho a^3} \right) + o(\rho^2 a (\rho a^3)^{1/2}).$$

- Lenz (1929), Bogoliubov (1947), Lee-Huang-Yang (1957).
- Rigorous proof of leading term Dyson (1957, upper), Lieb-Yngvason (1998).
- Upper bounds giving second order term: Erdős-Schlein-Yau (2008), Yau-Yin (2009), (Aaen 2014), Basti-Cenatiempo-Schlein (2021). Hard core is open but recent progress on hard core in GP-limit (Basti-Cenatiempo-Olgianti-Pasqualetti-Schlein).

The 3D Lee-Huang-Yang formula

Theorem

In the dilute limit $\rho a^3 \rightarrow 0$,

$$e^{3D}(\rho) = 4\pi\rho^2 a \left(1 + \frac{128}{15\sqrt{\pi}} \sqrt{\rho a^3} \right) + o(\rho^2 a (\rho a^3)^{1/2}).$$

- Lenz (1929), Bogoliubov (1947), Lee-Huang-Yang (1957).
- Rigorous proof of leading term Dyson (1957, upper), Lieb-Yngvason (1998).
- Upper bounds giving second order term: Erdős-Schlein-Yau (2008), Yau-Yin (2009), (Aaen 2014), Basti-Cenatiempo-Schlein (2021). Hard core is open but recent progress on hard core in GP-limit (Basti-Cenatiempo-Olgianti-Pasqualetti-Schlein).
- Lower bound in all cases SF-Solovej 2020-21.

The 2D formula

Theorem (SF, Girardot, Junge, Morin and Olivieri)

In the dilute limit $\rho a^2 \rightarrow 0$,

$$e^{2D}(\rho) = 4\pi\rho^2\delta_0 \left(1 + (2\Gamma + \frac{1}{2} + \log(\pi))\delta_0 \right) + o(\rho^2\delta_0^2),$$

with

$$\delta_0 := \frac{1}{|\log(\rho a^2 |\log(\rho a^2)|^{-1})|} = \frac{1}{|\log(\rho a^2)| + \log(|\log(\rho a^2)|)}$$

where $\Gamma = 0.577\dots$ is the Euler-Mascheroni constant.

- Schick (1971), Hines-Frankel-Mitchell (1978), Cherny-Shanenko (2001), Mora-Castin (2009), Yang (2008).

The 2D formula

Theorem (SF, Girardot, Junge, Morin and Olivieri)

In the dilute limit $\rho a^2 \rightarrow 0$,

$$e^{2D}(\rho) = 4\pi\rho^2\delta_0 \left(1 + (2\Gamma + \frac{1}{2} + \log(\pi))\delta_0 \right) + o(\rho^2\delta_0^2),$$

with

$$\delta_0 := \frac{1}{|\log(\rho a^2 | \log(\rho a^2)|^{-1})|} = \frac{1}{|\log(\rho a^2)| + \log(|\log(\rho a^2)|)}$$

where $\Gamma = 0.577\dots$ is the Euler-Mascheroni constant.

- Schick (1971), Hines-Frankel-Mitchell (1978), Cherny-Shanenko (2001), Mora-Castin (2009), Yang (2008).
- Leading order by Lieb-Yngvason (2001).

The 2D formula

Theorem (SF, Girardot, Junge, Morin and Olivieri)

In the dilute limit $\rho a^2 \rightarrow 0$,

$$e^{2D}(\rho) = 4\pi\rho^2\delta_0 \left(1 + (2\Gamma + \frac{1}{2} + \log(\pi))\delta_0 \right) + o(\rho^2\delta_0^2),$$

with

$$\delta_0 := \frac{1}{|\log(\rho a^2 | \log(\rho a^2)|^{-1})|} = \frac{1}{|\log(\rho a^2)| + \log(|\log(\rho a^2)|)}$$

where $\Gamma = 0.577\dots$ is the Euler-Mascheroni constant.

- Schick (1971), Hines-Frankel-Mitchell (1978), Cherny-Shanenko (2001), Mora-Castin (2009), Yang (2008).
- Leading order by Lieb-Yngvason (2001).
- GP regime Caraci-Cenatiempo-Schlein (2021-22).

The 2D formula

Theorem (SF, Girardot, Junge, Morin and Olivieri)

In the dilute limit $\rho a^2 \rightarrow 0$,

$$e^{2D}(\rho) = 4\pi\rho^2\delta_0 \left(1 + (2\Gamma + \frac{1}{2} + \log(\pi))\delta_0 \right) + o(\rho^2\delta_0^2),$$

with

$$\delta_0 := \frac{1}{|\log(\rho a^2 |\log(\rho a^2)|^{-1})|} = \frac{1}{|\log(\rho a^2)| + \log(|\log(\rho a^2)|)}$$

where $\Gamma = 0.577\dots$ is the Euler-Mascheroni constant.

- Schick (1971), Hines-Frankel-Mitchell (1978), Cherny-Shanenko (2001), Mora-Castin (2009), Yang (2008).
- Leading order by Lieb-Yngvason (2001).
- GP regime Caraci-Cenatiempo-Schlein (2021-22).
- Both upper and lower bound for general potentials including hard core.

2D is hard core - even for soft potentials

Using the constant function as trial state gives

$$e(\rho) \leq \frac{1}{2} \rho^2 \int v.$$

This is the wrong **constant** in $3D$ but the wrong **order** in $2D$.



2D is hard core - even for soft potentials

Using the constant function as trial state gives

$$e(\rho) \leq \frac{1}{2}\rho^2 \int v.$$

This is the wrong **constant** in 3D but the wrong **order** in 2D.

BEC

Our understanding of the energy is based on the Bogoliubov approximation that uses BEC as an input. But there is no BEC in 2D at low positive temperature.



2D is hard core - even for soft potentials

Using the constant function as trial state gives

$$e(\rho) \leq \frac{1}{2}\rho^2 \int v.$$

This is the wrong **constant** in $3D$ but the wrong **order** in $2D$.

BEC

Our understanding of the energy is based on the Bogoliubov approximation that uses BEC as an input. But there is no BEC in $2D$ at low positive temperature. However, we only use and prove BEC on finite length scales. On these scales there is (we can prove it!) BEC in both dimensions.



Upper bound: Reduction to large box

The central trial state corresponds to a **periodic** box of sidelength L^u .

Upper bound: Reduction to large box

The central trial state corresponds to a **periodic** box of sidelength L^u . The state has N particles and density $\rho = N/(L^u)^d$.

Upper bound: Reduction to large box

The central trial state corresponds to a **periodic** box of sidelength L^u . The state has N particles and density $\rho = N/(L^u)^d$.

This **periodic** state is then extended to a Dirichlet state of side length $L^u + \ell^u$. Finally, copies of these Dirichlet states are used to fill up the thermodynamic box.

Upper bound: Reduction to large box

The central trial state corresponds to a **periodic** box of sidelength L^u . The state has N particles and density $\rho = N/(L^u)^d$.

This **periodic** state is then extended to a Dirichlet state of side length $L^u + \ell^u$. Finally, copies of these Dirichlet states are used to fill up the thermodynamic box.

Localization error $N(\ell^u)^{-2} N(L^u \ell^u)^{-1}$,

New density $\tilde{\rho} = N/(L^u + \ell^u)^d \approx \rho(1 + (\ell^u/L^u))$.

Upper bound: Reduction to large box

The central trial state corresponds to a **periodic** box of sidelength L^u . The state has N particles and density $\rho = N/(L^u)^d$.

This **periodic** state is then extended to a Dirichlet state of side length $L^u + \ell^u$. Finally, copies of these Dirichlet states are used to fill up the thermodynamic box.

Localization error $N(\ell^u)^{-2} \approx N(L^u \ell^u)^{-1}$,

New density $\tilde{\rho} = N/(L^u + \ell^u)^d \approx \rho(1 + (\ell^u/L^u))$.

Need

$$\rho(L^u \ell^u)^{-1} \ll \begin{cases} a\rho^2 \sqrt{\rho a^3}, & d = 3, \\ \rho^2 / (\log(\rho a^2))^2, & d = 2, \end{cases} \quad \text{and} \quad \ell^u / L^u \ll \begin{cases} \sqrt{\rho a^3}, & d = 3, \\ (\log(\rho a^2))^{-1}, & d = 2. \end{cases}$$

Upper bound: Reduction to large box

The central trial state corresponds to a **periodic** box of sidelength L^u . The state has N particles and density $\rho = N/(L^u)^d$.

This **periodic** state is then extended to a Dirichlet state of side length $L^u + \ell^u$. Finally, copies of these Dirichlet states are used to fill up the thermodynamic box.

Localization error $N(\ell^u)^{-2} \quad N(L^u \ell^u)^{-1}$,

New density $\tilde{\rho} = N/(L^u + \ell^u)^d \approx \rho(1 + (\ell^u/L^u))$.

Need

$$\rho(L^u \ell^u)^{-1} \ll \begin{cases} a\rho^2 \sqrt{\rho a^3}, & d = 3, \\ \rho^2 / (\log(\rho a^2))^2, & d = 2, \end{cases} \quad \text{and} \quad \ell^u / L^u \ll \begin{cases} \sqrt{\rho a^3}, & d = 3, \\ (\log(\rho a^2))^{-1}, & d = 2. \end{cases}$$

So

$$L^u \gg a \begin{cases} (\rho a^3)^{-1}, & d = 3, \\ (\rho a^2)^{-1/2} (\log(\rho a^2))^{3/2}, & d = 2. \end{cases} \quad N = \rho(L^u)^d \gg \begin{cases} (\rho a^3)^{-2}, & d = 3, \\ (\log(\rho a^2))^3, & d = 2. \end{cases}$$

- Localize to boxes of size $\ell \gg \rho^{-\frac{1}{2}} |\log(\rho a^2)|^{\frac{1}{2}}$. Localization needs to preserve 'Neumann gap'. To get a priori information localize to smaller boxes of size $\ll (\rho)^{-\frac{1}{2}} (\log(\rho a^2))^{-\frac{1}{2}}$. Here Neumann gap can be used to control errors. Rest of analysis carried out on large box. The interaction between localized particles is denoted by $w(x_i, x_j)$.
- Condensation. Let P projection on constant function, Q orthogonal complement.

$$n_0 = \sum P_i, \quad n_+ = \sum Q_i.$$

A priori bounds control expected values $\langle n_0 \rangle$ and $\langle n_+ \rangle$. Energy error negligible if localizing to subspace where $n_+ \leq \mathcal{M}$ $n_+^L \leq \mathcal{M}$ where \mathcal{M} is between $\langle n_+ \rangle$ and n and where n_+^L counts the number of excitations with low momentum ($\lesssim \ell^{-1}$).



Thank you for your attention.

