

Dilute Bose and Fermi gases in dimensions 1–3

Jan Philip Solovej
Department of Mathematical Sciences
University of Copenhagen

Herrsching
June 9, 2022

Based on joint work with Søren Fournais and with
Johannes Agerkov and Robin Reuvers

Outline of Talk

- 1 Many-body Hamiltonians with 2-body interactions
- 2 Thermodynamic limit
- 3 Examples
- 4 Exactly solvable examples in 1D
- 5 The scattering length
- 6 The dilute limit
- 7 Dilute limit for fermions
- 8 The 1D upper bound
- 9 The 1D lower bound: Dyson's Lemma
- 10 The 1D lower bound: Reducing to the Lieb-Liniger model
- 11 A norm estimate
- 12 Conclusion

Many-body Hamiltonians with 2-body interactions

Consider N identical particles in a box $\Omega_L = [0, L]^d$ (Dirichlet or periodic b.c.). Dimension: $d = 1, 2, 3$.

The most basic translation invariant many-body Hamiltonian:

$$H_N = \sum_{i=1}^N -\Delta_i + \sum_{1 \leq i < j \leq N} v(x_i - x_j)$$

2-body potential: $v : \mathbb{R}^d \rightarrow [0, \infty]$ spherically symmetric, compact support.

Hilbert Space Bosons: $\mathcal{H}_N(\Omega_L) = \bigvee^N L^2(\Omega_L)$ or $L^2(\Omega_L^N)$

Hilbert Space (spin- J) Fermions: $\mathcal{H}_N(\Omega_L) = \bigwedge^N L^2(\Omega_L; \mathbb{C}^{2J+1})$

The thermodynamic limit

Our focus will be on the **Ground State Energy**:

$$E_L(N) = \inf \text{Spec}_{\mathcal{H}_N(\Omega_L)}(H_N)$$

We will be interested in the **Thermodynamic Limit**:

$$e(\rho) = \lim_{\substack{L \rightarrow \infty \\ N/L^d \rightarrow \rho}} e_L(N)$$

of the **energy per volume**

$$e_L(N) = L^{-d} E_L(N).$$

We will write $e_B(\rho)$ or $e_F(\rho)$ for Bosons or Fermions.

Examples

Free gas: $v = 0$

$$e_B^{\text{free}}(\rho) = 0, \quad e_F^{\text{free}}(\rho) = C_d \rho^{(d+2)/d}, \quad d = 1 : \quad e_F^{\text{free}}(\rho) = \frac{\pi^2}{3} \rho^3$$

The Bosonic state is **pure condensate**:

$$\Psi(x_1, \dots, x_N) = \prod_{i=1}^N \phi(x_i), \quad \phi \approx \text{const.}$$

If we use this as a trial state for $v \neq 0$ we get $e_B(\rho) \leq \frac{1}{2} \rho^2 \int v$.

Note for 1D:

$$e_F^{\text{free}}(\rho) \leq \frac{1}{2} \rho^2 \int v, \quad \text{if } \rho \leq \frac{3}{2\pi^2} \int v.$$

Hard core potential:

$$v_a(x) = \begin{cases} \infty, & |x| < a \\ 0, & |x| > a \end{cases}$$

Exactly solvable examples in 1D

For $d = 1$ we have for the **hard core gas**

$$e_B^{\text{HC}}(\rho) = e_F^{\text{HC}}(\rho) = \frac{\pi^2}{3} \rho^3 (1 - \rho a)^{-2}$$

Like a free Fermi gas $\frac{\pi^2}{3} N \rho^2$ with a smaller volume:
 $\rho \rightarrow \rho(1 - \rho a)^{-1}$.

The **Lieb-Liniger model** for bosons has $v = 2c\delta_0$. It can be solved exactly (Lieb-Liniger 1963). In particular,

$$e_B^{\text{LL}}(\rho) \geq \frac{\pi^2}{3} \rho^3 (1 - \rho a)^{-2}, \quad a = -2/c.$$

The lower bound is asymptotically right as $\rho|a| \rightarrow 0$.

Main goal here is to explain that $\frac{\pi^2}{3} \rho^3 (1 - \rho a)^{-2}$ is asymptotically correct for all v .

The scattering length

For $\text{supp}(v) \subseteq B_R$. Let ϕ be unique solution to

$$-\Delta\phi + \frac{1}{2}v\phi = 0, \quad \text{on } B_R, \quad \phi(x) = 1 \text{ for } |x| = R$$

Then $\phi(x) = f(|x|)$ and for $\text{range}(v) < r < R$, we have

$$f(r) = \begin{cases} (r - a)/(R - a) & \text{for } d = 1 \\ \ln(r/a)/\ln(R/a) & \text{for } d = 2 \\ (1 - ar^{2-d})/(1 - aR^{2-d}) & \text{for } d \geq 3, \end{cases}$$

with some constant $a \in \mathbb{R} \cup \{\infty\}$ called the (s-wave) **scattering length**.

For hard core and Lieb-Liniger a is the scattering length.

For $v = 0$, $\phi = 1$, i.e., $a = 0$ for $d \geq 2$, $a = \pm\infty$ for $d = 1$.

Dilute limit for bosons

Theorem (Dilute limit for bosons)

If range of v is R_0 then as $\rho|a|^d \rightarrow 0$

$$1D: e_B(\rho) = \frac{\pi^2}{3} \rho^3 \left[(1 - \rho a)^{-2} + O((\rho R)^{6/5}) \right].$$

with $R = \max\{R_0, 2|a|\}$. Error depends on $R \int v_{\text{reg}}$.

$$2D: e_B(\rho) = 4\pi\rho^2 \left(|\ln(\rho a^2)|^{-1} + o(|\ln(\rho a^2)|^{-1}) \right).$$

Error depends on R_0/a .

$$3D: e_B(\rho) = 4\pi\rho^2 a \left(1 + \frac{128}{15\sqrt{\pi}} \sqrt{\rho a^3} + o(\sqrt{\rho a^3}) \right).$$

Upper bound requires $v \in L^1$. Error depends on R_0/a .

The 1D case is joint work with Agerskov and Reuvers.

The 2D case is work of Lieb-Yngvason.

The lower bound in 3D is joint work Fournais.

The upper bound in 3D is Yau-Yin and Basti-Cenatiempo-Schlein.

Dilute limit for fermions

Theorem (Spin-1/2 fermions in 3D, Lieb-Seiringer-Solovej)

$$e_F(\rho_\uparrow, \rho_\downarrow) = C_3(\rho_\uparrow^{5/3} + \rho_\downarrow^{5/3}) + 8\pi a \rho_\uparrow \rho_\downarrow + o(a\rho^2)$$

Theorem (Spin-less fermions in 1D, Agerskov-Reuvers-Solovej)

$$e_F(\rho) = \frac{\pi^2}{3} \rho^3 \left[(1 - \rho a_p)^{-2} + O(\rho a_p)^{6/5} \right].$$

Here $a_p \geq 0$ is the 1D p -wave scattering length.

Dilute limit for fermions (Continued)

Theorem (Spin-1/2 fermions in 1D, Agerskov-Reuvers-Solovej 2022)

The absolute (total spin zero) ground state energy density satisfies

$$e_F(\rho) \leq \frac{\pi^2}{3} \rho^3 [1 - 2e_H \rho a + 2(1 + e_H) \rho a_p + o(\rho a_p)].$$

where e_H is the thermodynamic limit of the ground state energy per spin of the Heisenberg antiferromagnet chain with Hamiltonian

$$\sum_i S_i \cdot S_{i+1}.$$

In fact, $e_H = -\ln 2$ (Bethe-Hulthén). We **conjecture** that this is the correct asymptotics. The result agrees with the hard core case where $a = a_p$ and the delta-function gas ($a_p = 0$) (Yang).

The 1D upper bound (bosons)

The trial state has to capture free Fermi energy, as well as correction due to scattering process. Hence we consider for $x = (x_1, \dots, x_N)$

$$\Psi(x) = \begin{cases} \omega(\mathcal{R}(x)) \frac{|\Psi_F(x)|}{\mathcal{R}(x)} & \text{if } \mathcal{R}(x) < b \\ |\Psi_F(x)| & \text{if } \mathcal{R}(x) \geq b \end{cases}, \quad \mathcal{R}(x) = \min_{i < j} (|x_i - x_j|)$$

ω is the suitably normalized ($\omega(b) = b$) **scattering solution** and Ψ_F is **the free Fermi gas** wave function (Dirichlet b.c.)

We can calculate higher correlation functions for Ψ_F using Wick's Theorem. We use the diamagnetic "inequality"

$$\int |\nabla |\Psi_F||^2 \leq \int |\nabla \Psi_F|^2.$$

Localization: We get errors that behave badly in N and we must therefore localize, i.e. write the trial state as products from smaller boxes.

The 1D lower bound: Dyson's Lemma

Lemma (Dyson)

Let $R > R_0 = \text{range}(v)$ and $\varphi \in H^1(\mathbb{R})$, then for any interval $\mathcal{I} \ni 0$

$$\int_{\mathcal{I}} |\partial\varphi|^2 + \frac{1}{2}v|\varphi|^2 \geq \int_{\mathcal{I}} \frac{2}{R-a} (\delta_R + \delta_{-R}) |\varphi|^2,$$

where a is the s -wave scattering length.

Hence for a many-body wave function we have, denoting

$$\mathbf{r}_i(x) = \min_{j \neq i} (|x_i - x_j|)$$

$$\int \sum_i |\partial_i \Psi|^2 + \sum_{i < j} v(x_i - x_j) |\Psi|^2 \geq \int \sum_i |\partial_i \Psi|^2 \chi_{\mathbf{r}_i(x) > R} + \sum_i \frac{2}{R-a} \delta(\mathbf{r}_i(x) - R) |\Psi|^2.$$

The 1D lower bound: Reducing to the Lieb-Liniger model

Define $\psi \in L^2([0, \ell - (n-1)R]^n)$ by

$$\psi(x_1, x_2, \dots, x_n) = \Psi(x_1, R + x_2, \dots, (n-1)R + x_n),$$

for $x_1 \leq x_2 \leq \dots \leq x_n$ and symmetrically extended. Then (ignoring some finite volume and lower order corrections)

$$\begin{aligned} \langle \Psi | H_N | \Psi \rangle &\geq E_{LL}(N, \tilde{L} = L - NR, \tilde{c} = 2/(R - a)) \langle \psi | \psi \rangle \\ &\geq N \frac{\pi^2}{3} \rho^2 (1 - \rho R)^{-2} (1 + \rho(R - a))^{-2} \langle \psi | \psi \rangle \\ &\geq N \frac{\pi^2}{3} \rho^2 (1 + 2\rho R - 2\rho(R - a)) \langle \psi | \psi \rangle \\ &= N \frac{\pi^2}{3} \rho^2 (1 + 2\rho a) \langle \psi | \psi \rangle. \end{aligned}$$

We need to control that $\langle \psi | \psi \rangle$ is close to 1.

A norm estimate

$$1 - \langle \psi | \psi \rangle \leq \text{const.} \cdot R^2 \left(\sum_{i < j} \int_{B_{ij}} |\partial_i \Psi|^2 + \sum_{i < j} \int v_{ij} |\Psi|^2 \right).$$

where $B_{ij} = \{x \mid \tau_i(x) < R\}$ and we assume $R > \max\{R_0, 2|a|\}$. This allows us to control the missing mass by the energy using a bootstrap argument. Again there are errors that behave badly in N and we must localize by going to a **grand canonical formulation** (choose μ appropriately):

$$\begin{aligned} E_L(N) &\geq M \inf_n (E_{L/M}^{\text{Neumann}}(n) - \mu n) + \mu N \\ &\geq M \inf_{n \leq (\text{const.})\rho L/M} (E_{L/M}^{\text{Neumann}}(n) - \mu n) + \mu N. \end{aligned}$$

Now the errors will depend only on $\rho L/M$ and this will close the argument.

Conclusion

I have discussed the asymptotics of the ground state energy of dilute Bose and also Fermi gases in 1–3 dimensions.

Some open problems are:

- General Spin-1/2 Fermions in 1–3 D
- Higher order corrections
- Large density limits