

Some Mathematical Questions of Quantum Mechanics

I.M.Sigal
Toronto

Work with V. Bach, J. Fröhlich, W. Hunziker, A. Soffer,
Th. Chen, J. Faupin, Zhou Gang, M. Griesemer, S. Gustafson,
S. Breteaux, I. Chenn, L. Jonsson, F. Pusateri, T. Tzaneteas
Discussions: R. Frank, G.M. Graf, Ch. Hainzl and B. Pausader

Herrsching, June 2022

Schrödinger equation

Many-body quantum systems are described by the [Schrödinger eq.](#)

$$i\partial_t\psi = H_n\psi. \quad (\text{SE})$$

where $\psi = \psi(x_1, \dots, x_n, t)$ and $H_n = n$ -particle Schrödinger opr,

$$H_n := \sum_1^n \frac{-1}{2m} \Delta_{x_i} + \sum_{i < j} v(x_i - x_j). \quad (1)$$

(For n particles of mass m interacting via a 2-body potential v .)

Global existence & unitar. \iff [self-adjointness](#) of H_n

Goal: Describe the space-time behaviour of solutions

Main problem: *stability* vs *break-up* or *collapse*.

Stability = localiz. in space & period. in time (atoms, ..., stars):

- ▶ *stability w. r. to collapse* ($\inf H_n > -\infty$)
- ▶ *stability w. r. to break-up* ($\inf H_n =$ [eigenvalue](#)).

Break-up (Local decay) \implies scattering

Scattering

The main mathematical problem of the scattering theory

Asymptotic completeness: As time progresses, a quantum system settles in a superposition of states in each of which it is broken into a stable freely moving fragments.

Theorem (Soffer-IMS ($\mu > 1$), Dereziński ($\mu > \sqrt{3} - 1$))

Suppose that the pair potentials $v_{ij}(x_i - x_j)$ entering H_n satisfy $v_{ij}(y) = O(|y|^{-\mu})$, with $\mu > \sqrt{3} - 1$. Then the asymptotic completeness holds.

Earlier works: [Deift](#), [Enss](#), [Gérard](#), [Graf](#), [Mourre](#), [Simon](#), [Yafaev](#), ...

Open problem: Prove the asymptotic completeness $v_{ij}(y) = O(|y|^{-\mu})$, with $\mu \leq \sqrt{3} - 1$.

Including photons (NR QED)

To describe the real (at least visible) world, have to couple the particles to **photons** (quantized electromagnetic field) \implies

$$i\partial_t\psi_t = H_\kappa\psi_t,$$

where H_κ is the Hamiltonian on the state space $\mathcal{H} := \mathcal{H}_p \otimes \mathcal{H}_f$:

$$H_\kappa = \sum_{j=1}^n \frac{1}{2m} (-i\nabla_{x_j} - \kappa A_\xi(x_j))^2 + U(x) + H_f. \quad (2)$$

Here, $\kappa =$ particle charge, $U(x) =$ total potential, $H_f =$ photon Hamilt. n.

$A_\xi(y) =$ UV-regularized quantized vector potential.

Infrared problem: photons \implies infinite clouds of gapless 'excitations'.

Main qstn: Math. descript. of the processes of emiss. and absorp. of rad.

Bach-Fröhlich-IMS: stability of the *ground state* and instability of the *excited states* of the particle system (say, an atom) and emergence of resonances.

Further results: Th. Chen, Griesemer-Lieb-Loss, Hasler-Herbst, Hiroshima, Hainzl-Seiringer, Hübner-Spohn, Miyao, Møller, A. Panati, Pizzo, Teufel, et al.

Scattering

Consider the **Rayleigh scattering**, i.e. the scattering at the **energy** $<$ **the ionization energy** of atomic or molecular system.

Thm (**Faupin-IMS, De Roeck-Griesemer-Kupiainen**). Assume that

$$\langle \psi_t, N_{\text{ph}} \psi_t \rangle \leq C < \infty \quad (\text{satisfied in spec. cases}^1).$$

Then the asymptotic completeness holds.

Earlier results: Spohn, Dereziński-Gérard, Fröhlich-Griesemer-Schlein, et al

Open problem: Prove $\langle \psi_t, N_{\text{ph}} \psi_t \rangle \leq C < \infty$ for general particle systems (like atoms).

¹De Roeck-Kupiainen, spin-boson model, Faupin-IMS, some generalizations

Effective (Hartree and Hartree-Fock) Equations

Effective equations is a powerful tool in study of complex systems. For the n -particle quantum systems, a key effective equation is the **Hartree-Fock equation**

$$i\partial_t\gamma = [h_\gamma, \gamma], \quad (\text{HF})$$

where γ is a (trace class) positive operator (**density opr**) and h_γ is the **self-consistent** Hamiltonian:

$$h_\gamma := -\Delta + V + g(\gamma),$$

with the **self-interaction** energy $g(\gamma) = v \star \rho_\gamma + \text{ex}_{\text{HF}}(\gamma)$. Here

$$\rho_\gamma(x, t) := \gamma(x; x, t) \quad (\text{charge density}).$$

For fermions $\gamma \leq \mathbf{1}$, for bosons, $\gamma < \infty$ and $\text{ex}_{\text{HF}} = 0$

The HF eq **trades the degrees of freedom for nonlinearity**. It is widely used in quantum chemistry and condensed matter physics.

Density functional theory

Replacing the the exchange term, $ex_{\text{HF}}(\gamma)$, in the HF eq by a function $xc(\rho_\gamma)$ (exchange and correlation energy) of the 1-particle (charge) density ρ_γ (LDA), one arrives at the (t -dep) Kohn-Sham equation

$$i\partial_t\gamma = [h_{\rho_\gamma}, \gamma], \quad (\text{KS})$$

$$h_\rho := -\Delta + v * \rho + xc(\rho) \quad (\text{self-consist ham}), \quad (\text{SCH})$$

$$\rho_\gamma(x, t) := \gamma(x; x, t) \quad (\text{charge density}). \quad (\text{CD})$$

Density functional theory (DFT): Runge-Gross (modelled on Hohenberg-Kohn and Levy-Lieb theories): under certain conditions,

$\exists xc(\rho) : \quad i\partial_t\Psi = H_n\Psi \implies \gamma := \text{Tr}_{(n-1)} P_\Psi$ satisfies (KS).

Quant Chem./Cond. Matt. Phys. : Design $xc(\rho)$ depending on model.

Pusateri-IMS: Short-range scatt. $|xc(\rho)| \lesssim \rho^\beta, \beta > 1/\min(d, 2)$

Open problem: Long-range scattering

Earlier results: Reduced HF eq ($xc = 0$): Stability of ground states (Lewin-Sabin, Th. Chen-Pavlovic)

Quantum liquids

However, the H and HF equations fail to describe quantum matter at nano- and macro-scales: superconductors, superfluids and BE condensates. For this, one needs **another conceptual step**:

Pass to the most general quasi-free (one-body) approximation to the n -body dynamics, of which the HF approximation is a special case.

Such a step was made by Bardeen-Cooper-Schrieffer for fermions and by Bogolubov, for bosons, and resulted in²

⇒ The (time-dependent) *Bogolubov-de Gennes* (fermions)
and *Hartree-Fock-Bogolubov* (bosons) equations.

* * *

Key problem: Existence and stability of the ground/equilibrium state - static solution minimizing the free energy (locally, for translation invariance) ⇒ symmetry breaking.

²See Bach-Bret-Chen-Fröhl-IMS, Chenn-IMS, Benedicter-Sok-Solovej ▶

Hartree-Fock-Bogolubov system³

Neglecting the α -component and taking $v = \lambda\delta$, $\lambda \in \mathbb{R}$ for the pair interaction potential, the **HFB syst** becomes (2-gas model)

$$i\partial_t\phi = h\phi + \lambda|\phi|^2\phi + 2\lambda\rho_\gamma\phi, \quad (\text{GP})$$

$$i\partial_t\gamma = [h_{\gamma,\phi}, \gamma], \quad (\text{HF})$$

where $h = -\Delta + V$ is a one-part. Schrödinger opr, $\rho_\gamma(x, t) := \gamma(x; x, t)$,

$$h_{\gamma,\phi} := h + 2\lambda(\rho_\gamma + |\phi|^2). \quad (\text{scSO})$$

These are coupled Gross-Pitaevskii and Hartree-Fock equations

Here the condensate of atoms (ϕ) is coupled to a cloud of the thermal atoms (γ).

Problems: Existence, Ground (=equil) st., Condensation, Collapse oscillat. for $\lambda < 0$ (**correction to the Papanicolaou-Sulem² collapse law?**).

Partial results of transl. invar. case: [NapiorkowskiReuvSol](#), [BBCF](#)

³See Bach-Breteaue-Chen-Fröhlich-IMS

Bogolubov-de Gennes system

For **fermions** (electrons), (γ, α) couple to the **EM field**. Let a be the **magnetic** potential and take the gauge $\phi_{\text{electr}} = 0$. Then⁴

$$\begin{aligned}i\partial_t\gamma &= [h_{a,\gamma}, \gamma] + \dots, \\i\partial_t\alpha &= [h_{a,\gamma}, \alpha]_+ + \dots, \\-\partial_t^2 a &= \text{curl}^* \text{curl } a - j(\gamma, a),\end{aligned}\tag{BdG}$$

where $j(\gamma, a)(x) := [-i\nabla_a, \gamma]_+(x, x)$, the current density,

$$h_{a,\gamma} = -\Delta_a + g_{\text{xc}}(\gamma),\tag{3}$$

with $\Delta_a := (\nabla + ia)^2$, and $[A, B]_+ = AB + B\bar{A}$, $\bar{A} := CAC$.

These are the celebrated **Bogolubov-de Gennes equations**. They give the 'mean-field' (BCS) theory of superconductivity.

Key problem: Existence of the ground/equilibrium state (static soln minimizing the free energy locally) and **symmetry breaking?**

⁴Chenn-IMS, Benedicter-Sok-Solovej

Gauge (magnetic) translational invariance

For the ground state (GS), look for the most symmetric state(s).

The BdG eqs are invariant under the (t -indep.) *gauge* transforms

$$T_{\chi}^{\text{gauge}} : (\gamma, \alpha, \mathbf{a}) \rightarrow (e^{i\chi}\gamma e^{-i\chi}, e^{i\chi}\alpha e^{i\chi}, \mathbf{a} + \nabla\chi) \quad (4)$$

The simplest class of states: translationally invariant states for $\mathbf{a} = 0$ and the gauge-translationally invariant ones for $\mathbf{a} \neq 0$.

Gauge (magnetically) transl. invariant states are invariant under

$$T_{bs} : (\gamma, \alpha, \mathbf{a}) \rightarrow (T_{\chi_s}^{\text{gauge}})^{-1} T_s^{\text{transl}}(\gamma, \alpha, \mathbf{a}),$$

for any $s \in \mathbb{R}^d$. For $d = 2, 3$, $\chi_s(x) := \frac{b}{2} \cdot (s \wedge x)$ (special gauge).

Here T_s^{transl} , $s \in \mathbb{R}^d$, is the group of translations and $b > 0$ is a parameter identified with a *constant external magnetic field*.

Ground State

Usually, the **ground state (GS)** has the max. symmetry \Rightarrow

Depending on the magnetic field b , one expects:

GS is **translationally invariant** for $b = 0$,

GS is **magnetically translationally (MT) invariant** for $b \neq 0$.

Candidates for the ground state:

1. **Normal states**: (γ, α, a) , with $\alpha = 0$ ($\Rightarrow \gamma$ is Fermi-Dirac state).
2. **Superconducting states**: (γ, α, a) , with $\alpha \neq 0$ and $a = 0$.

Theorem (Hainzl-Hamza-Seiringer-Solovej). For $b = 0$, \exists superconducting, normal, **translationally invariant** solution.

Theorem (Chenn-IMS). For $b \neq 0$, MT-invariance \implies normality ($\alpha = 0$).

Corollary. For $b \neq 0$, superconductivity \implies symmetry breaking.

Vortex lattices

For the GS, look for states with minimal symmetry breaking \implies

► **Vortex lattice:** \exists lattice \mathcal{L} in \mathbb{R}^2 and map $\chi_s : \mathcal{L} \times \mathbb{R}^2 \rightarrow \mathbb{R}$:

$$T_s^{\text{transl}}(\gamma, \alpha, a) = T_{\chi_s}^{\text{gauge}}(\gamma, \alpha, a), \forall s \in \mathcal{L}, \text{ and } \alpha \neq 0.$$

T_s^{transl} is a group representation $\implies \chi_s$ satisfies the **co-cycle relat:**

$$\chi_{s+t}(x) - \chi_s(x+t) - \chi_t(x) = \frac{b}{2} \cdot (s \wedge t) \in 2\pi\mathbb{Z}, \forall s, t \in \mathcal{L}. \quad (5)$$

Co-cycle relation (5) \implies the **magnetic flux is quantized:**

$$\frac{1}{2\pi} \int_{\Omega^{\mathcal{L}}} \text{curl } a \equiv c_1(\chi) \in \mathbb{Z}.$$

Here $\Omega^{\mathcal{L}}$ is a fundamental cell of \mathcal{L} and $c_1(\chi)$ is the 1st Chern #.

Existence of vortex lattices

Theorem. For the BdG syst without the self-interact. term:

(i) $\forall n, T > 0$ and $\mathcal{L} \ni$ a static solution $u_{Tn\mathcal{L}} := (\gamma, \alpha, a)$ satisfying

$$u_{Tn\mathcal{L}} \text{ is } \mathcal{L}\text{-equivariant: } T_s^{\text{transl}} u_{Tn\mathcal{L}} = T_{\chi_s}^{\text{gauge}} u_{Tn\mathcal{L}}, \forall s \in \mathcal{L}, \quad (6)$$

$$\text{1st Chern number is } n: \int_{\Omega^{\mathcal{L}}} \text{curl } a = 2\pi n, \quad (7)$$

$u_{nT\mathcal{L}}$ minimizes the **free energy** $F_T = E - TS$ on $\Omega^{\mathcal{L}}$ for $c_1 = n$;

(ii) For the pair potential $v \leq 0, v \neq 0$ and T and b sufficiently small, $u_{Tn\mathcal{L}}$ is a **vortex lattice** (i.e. $\alpha \neq 0$);

(iii) For $n > 1$, there is a **finer lattice**, $\mathcal{L}' \supset \mathcal{L}$ for which $u_{Tn\mathcal{L}} = u_{T1\mathcal{L}'}$, i.e. $u_{Tn\mathcal{L}}$ is \mathcal{L}' -equivariant with $c_1 = 1$.


Remark: (ii) \implies **Symmetry breaking** in the ground state

Open problem: Is the vortex lattice a ground state?

Global minimizer (locally) vs **emerging local minimizer**.

Experiment

Experiment: the ground state is the hexagonal vortex lattice.



Theoretical description: the BdG system with the coarse-scale approximation given by the [Ginzburg-Landau system](#)

$$\begin{aligned}\Delta_a \psi + \kappa^2(1 - |\psi|^2)\psi &= 0, \\ -\text{curl}^2 a + \text{Im}(\bar{\psi} \nabla_a \psi) &= 0.\end{aligned}$$


($|\psi|^2$ is the density of superconducting electrons)

Ginzburg-Landau equations: vortex lattices

Thm: For type II superconductors ($\kappa > 1/\sqrt{2}$), $\exists b_*$ s.t. for $b > b_*$, the homogeneous soln, u_b^{hom} , with the constant magnetic field b , is a local minimizer of $E_{\text{GL}}(u)$ and for $b < b_*$, it is not.


$\forall b < b_*$ and lattice \mathcal{L} , \exists a soln, $u_b^{\mathcal{L}}$, with the lattice \mathcal{L} symmetry, with

$$\frac{1}{2\pi} \int_{\Omega^{\mathcal{L}}} \text{curl } a = b, \quad \text{and} \quad E_{\text{GL}}(u^{\mathcal{L}}) < E_{\text{GL}}(u^{\text{hom}}).$$



Thm (Tzanet-IMS): Soln $u_b^{\mathcal{L}}$ is formed by magn. vortices

arranged in a lattice \mathcal{L} .



Thm (Tzanet-IMS): Vort. lattices are local minima of the GL energy w.r.to (i.e. stable under) period & local perts.

Problem: Stability under more general lattice deformations?

Discussion: Global minimizer (locally) vs bifurcation of local minimizer?

Summary

We reviewed some results on quantum many-particle problem:

- ▶ n -particle scattering
- ▶ interaction of radiation and matter
- ▶ effective equations for quantum gases and quantum liquids (HF, KS-DFT, HFB, BdG, GL equations)

I highlighted a number of basic open problems in these central areas of quantum mechanics. They concentrated on

- ▶ Long-time behaviour
- ▶ Existence and stability of ground states
- ▶ Symmetry breaking

Thank-you for your attention